

ON δ_m CONSTANT LOCUS OF VERSAL DEFORMATIONS OF NONDEGENERATE HYPERSURFACE SIMPLE $K3$ SINGULARITIES

MASAKO FURUYA

INTRODUCTION

Simple $K3$ singularities are regarded as natural generalizations in three-dimensional case of simple elliptic singularities. The notion of a simple $K3$ singularity was defined by S. Ishii and K. Watanabe [IW] as a three-dimensional Gorenstein purely elliptic singularity of $(0,2)$ -type, whereas a simple elliptic singularity is a two-dimensional purely elliptic singularity of $(0,1)$ -type. It is also pointed out in [IW] that a simple $K3$ singularity is characterized as a quasi-Gorenstein singularity such that the exceptional set of any minimal resolution is a normal $K3$ surface. Let $f \in \mathbf{C}[x, y, z, w]$ be a polynomial which is nondegenerate with respect to its Newton boundary $\Gamma(f)$ in the sense of [V1], and whose zero locus $X = \{f = 0\}$ in $\mathbf{C}^4$ has an isolated singularity at the origin $0 \in \mathbf{C}^4$. Then the condition for $(X, 0)$ to be a simple $K3$ singularity is given by a property of the Newton boundary $\Gamma(f)$ of f (cf. Theorem 1.6).

Hypersurface simple $K3$ singularities defined by nondegenerate quasi-homogeneous polynomials are classified into ninety five classes in term of the weight of the polynomial by Yonemura [Yo]. We consider versal deformations of them. It has been conjectured that the stratum $\mu = \text{const}$ of the versal deformation of any nondegenerate hypersurface simple $K3$ singularity is equivalent to the δ_m constant locus by Ishii. It holds true for the case deformations are also nondegenerate by 1.7 (1) [W]. On the other hand, it follows from 2.2 ([R1], [R2]) that the δ_m constant locus includes the μ constant locus generally. We show the conjecture holds true in general for No.10-14, 46-51 and 83 in the table of [Yo].

I would like to express my sincere gratitude to Professor Shihoko Ishii for telling me about the conjecture and giving valuable advice. I also express my gratitude to Professors Masataka Tomari and Kei-ichi Watanabe for their useful comments concerning Theorem 2.4, and to Professor Takao Fujita for his helpful remark on

Section 1 and 2. I also thank Mr. Hironobu Ishihara who pointed out grammatical mistakes throughout this paper.

1. PRELIMINARY

In this section, we recall some definitions and facts from [I1], [IW], [W] and [Yo].

First we define the plurigenera δ_m , $m \in \mathbf{N}$, for normal isolated singularities and define purely elliptic singularities. Let (X, x) be a normal isolated singularity in an n -dimensional analytic space X , and $\pi : (\tilde{X}, E) \longrightarrow (X, x)$ a good resolution. In the following, we assume that X is a sufficiently small Stein neighbourhood of x .

Definition 1.1 (Watanabe [W]-Def. 1.2). Let (X, x) be a normal isolated singularity. For any positive integer m ,

$$\delta_m(X, x) := \dim_{\mathbf{C}} \Gamma(X - \{x\}, \mathcal{O}(mK)) / L^{2/m}(X - \{x\}),$$

where K is the canonical line bundle on $X - \{x\}$, and $L^{2/m}(X - \{x\})$ is the set of all $L^{2/m}$ -integrable (at x) holomorphic m -ple n -forms on $X - \{x\}$.

Then δ_m is finite and does not depend on the choice of a Stein neighbourhood X .

Definition 1.2 (Watanabe [W]-Def. 3.1). A singularity (X, x) is said to be purely elliptic if $\delta_m(X, x) = 1$ for every $m \in \mathbf{N}$.

In the following, we assume that (X, x) is quasi-Gorenstein, i.e., there exists a non-vanishing holomorphic 3-form on $X - \{x\}$. Let $E = \bigcup E_i$ be the decomposition of the exceptional set E into irreducible components, and write $K_{\tilde{X}} = \pi^* K_X + \sum_{i \in I} m_i E_i - \sum_{j \in J} m_j E_j$ with $m_i \geq 0$, $m_j > 0$. Ishii [I1] defined the essential part of the exceptional set E as $E_J = \sum_{j \in J} m_j E_j$, and showed that if (X, x) is purely elliptic, then $m_j = 1$ for all $j \in J$.

Definition 1.3 (Ishii [I1]-Def. 4.1). A quasi-Gorenstein purely elliptic singularity (X, x) is of $(0, i)$ -type if $H^{n-1}(E_J, \mathcal{O}_{E_J})$ consists of the $(0, i)$ -Hodge component $H_{n-1}^{0,i}(E_J)$, where

$$\mathbf{C} \cong H^{n-1}(E_J, \mathcal{O}_{E_J}) = Gr_F^0 H^{n-1}(E_J) = \bigoplus_{i=0}^{n-1} H_{n-1}^{0,i}(E_J).$$

Definition-Proposition 1.4 (Ishii-Watanabe [IW]-Def. 4). *A three-dimensional singularity (X, x) is a simple K3 singularity if the following two equivalent conditions are satisfied:*

- (1) *(X, x) is Gorenstein purely elliptic of $(0, 2)$ -type.*
- (2) *The exceptional divisor E is a normal K3 surface for any minimal resolution $\pi : (\tilde{X}, E) \longrightarrow (X, x)$.*

Recall 1.5. A minimal resolution $\pi : (\tilde{X}, E) \longrightarrow (X, x)$ is a proper morphism with $\tilde{X} - E \cong X - \{x\}$, where $\tilde{X}$ has only terminal $\mathbf{Q}$ -factorial singularities and $K_{\tilde{X}}$ is numerically effective with respect to π .

Next we consider the case where (X, x) is a hypersurface singularity defined by a nondegenerate polynomial $f = \sum a_\nu z^\nu \in \mathbf{C}[z_0, z_1, \dots, z_n]$, and $x = 0 \in \mathbf{C}^{n+1}$.

Recall 1.6. The Newton boundary $\Gamma(f)$ of f is the union of the compact faces of $\Gamma_+(f)$, where $\Gamma_+(f)$ is the convex hull of $\bigcup_{a_\nu \neq 0} (\nu + \mathbf{R}_{\geq 0}^{n+1})$ in $\mathbf{R}^{n+1}$. For any face Δ of $\Gamma_+(f)$, set $f_\Delta := \sum_{\nu \in \Delta} a_\nu z^\nu$. We say f to be nondegenerate, if

$$\frac{\partial f_\Delta}{\partial z_0} = \frac{\partial f_\Delta}{\partial z_1} = \dots = \frac{\partial f_\Delta}{\partial z_n} = 0$$

has no solution in $(\mathbf{C} - \{0\})^{n+1}$ for any face Δ .

When f is nondegenerate, the condition for (X, x) to be a purely elliptic singularity is given as follows:

Theorem 1.7 (Watanabe [W]-Prop. 2.9, Cor. 3.14). *Let f be a nondegenerate polynomial and suppose $X = \{f = 0\}$ has an isolated singularity at $x = 0 \in \mathbf{C}^{n+1}$.*

- (1) *(X, x) is purely elliptic if and only if $(1, 1, \dots, 1) \in \Gamma(f)$.*
- (2) *Let $n=3$ and let Δ_0 be the face of $\Gamma(f)$ containing $(1, 1, 1, 1)$ in the relative interior of Δ_0 . Then (X, x) is a simple K3 singularity if and only if $\dim_{\mathbf{R}} \Delta_0 = 3$.*

Thus if f is nondegenerate and defines a simple K3 singularity, then $f_{\Delta_0} := \sum_{\nu \in \Delta_0} a_\nu z^\nu$ is a quasi-homogeneous polynomial of a uniquely determined weight α called the weight of f and denoted $\alpha(f)$. Namely, $\alpha = (\alpha_1, \dots, \alpha_4) \in \mathbf{Q}_{>0}^4$ and $\deg_\alpha(\nu) := \sum_{i=1}^4 \alpha_i \nu_i = 1$ for any $\nu \in \Delta_0$. In particular, $\sum_{i=1}^4 \alpha_i = 1$, since $(1, 1, 1, 1)$ is always contained in Δ_0 .

Theorem 1.8 (Yonemura [Yo]-Prop. 2.1). *The cardinality of $\{\alpha(f) ; f \text{ is nondegenerate and defines a simple } K3 \text{ singularity, } \alpha_1 \geq \alpha_2 \geq \alpha_3 \geq \alpha_4\}$ is 95.*

In Table 2.2 of [Yo] can be found the complete list of weights $\alpha = \alpha(f)$ and examples of $f = \sum \alpha_\nu z^\nu$ such that f is quasi-homogeneous and that $\{f = 0\} \subset \mathbf{C}^4$ has a simple $K3$ singularity at the origin $0 \in \mathbf{C}^4$.

We describe a weight $\alpha = \alpha(f)$ as $\alpha = (p_1/p, p_2/p, p_3/p, p_4/p)$, where p, p_i are positive integers with $\gcd(p_1, p_2, p_3, p_4) = 1$.

Next we consider the versal deformation of simple $K3$ singularity $(X = \{f = 0\}, 0)$ such that f is a nondegenerate quasi-homogeneous polynomial.

Definition 1.9. A deformation of an isolated singularity (X, x) is a flat family of singularities:

$$\mathfrak{X} = \{(X_t, x_t) ; t \in U \subset \mathbf{C}^N\} \xrightarrow{\text{flat}} U$$

such that $(X_0, x_0) \cong (X, x)$ as germs of holomorphic functions.

Where U is a sufficiently small open neighbourhood of 0 in $\mathbf{C}^N$.

Definition 1.10. A deformation $\mathfrak{X}$ of an isolated singularity (X, x) is versal if for any deformation $\mathfrak{X}' \longrightarrow U'$ of (X, x) , the following is satisfied:

$$\begin{array}{ccc} \mathfrak{X} \times_U U' & \xrightarrow{\text{isom}} & \mathfrak{X}' \longrightarrow U' \\ & & \downarrow \exists \text{ holomorphic} \\ \mathfrak{X} & \longrightarrow & U \end{array}$$

Theorem 1.11 ([KS], [Tj]). *The versal deformation $\mathfrak{X}$ of an isolated singularity $(\{f = 0\}, 0)$ is described by*

$$\mathfrak{X} = \{(\{f + \sum \lambda_i g_i = 0\}, 0) ; (\lambda_i) \in U \subset \mathbf{C}^N\},$$

where the g_i determine a $\mathbf{C}$ -basis of the vector space $\mathbf{C}\{z_0, \dots, z_n\}/(f, f_{z_0}, \dots, f_{z_n})$.

The purpose of this paper is to show that the following conjecture holds true for No.10-14, 46-51 and 83 in Table 2.2 of [Yo].

Problem 1.12 (Ishii). Let $(X = \{f = 0\}, 0)$ be a hypersurface simple $K3$ singularity defined by a nondegenerate quasi-homogeneous polynomial f , and let $\mathfrak{X}$ be the versal deformation of $(X, 0)$. Then,

$$\begin{aligned} & \{\lambda \in U \subset \mathbf{C}^N ; \mu(X, 0) = \mu(X_\lambda, 0), (X_\lambda, 0) \in \mathfrak{X}\} \\ &= \{\lambda \in U \subset \mathbf{C}^N ; 1 = \delta_m(X, 0) = \delta_m(X_\lambda, 0) \text{ for all } m \geq 1, (X_\lambda, 0) \in \mathfrak{X}\}. \quad \square \end{aligned}$$

Since $(X, 0)$ and $(X_\lambda, 0)$ are hypersurface isolated singularities, they are normal Gorenstein, and so

$$P_g = 0 \iff \delta_m = 0 \text{ for all } m \geq 1.$$

On the other hand, $(X_\lambda, 0)$ is a deformation of a purely elliptic singularity $(X, 0)$, so it is either rational or purely elliptic. Therefore, this problem is equivalent to:

$$\mu(X, 0) = \mu(X_\lambda, 0) \iff 1 = P_g(X, 0) = P_g(X_\lambda, 0).$$

Furthermore, since μ and P_g are upper semi-continuous in respect of deformation ([M], [Te], [E]-Thm. 1, [Ya]-Thm. 2.6), it is equivalent to:

$$\mu(X, 0) > \mu(X_\lambda, 0) \iff 1 = P_g(X, 0) > P_g(X_\lambda, 0) = 0.$$

2. REDUCTION OF THE PROBLEM

Considering some facts, we can reduce the problem posed in section 1 to one about the weight of the defining polynomial of a hypersurface singularity.

Theorem 2.1 (Varchenko [V2]-Thm. 2). *Let $f \in \mathbf{C}[z_0, \dots, z_n]$ be quasi-homogeneous of weight α with $\alpha_0 + \dots + \alpha_n = 1$, and $\{f = 0\}$ has an isolated singularity at $0 \in \mathbf{C}^{n+1}$. Let*

$$\mu := \mu(f, 0) < +\infty, \quad f_\lambda := f + \sum_{i=1}^{\mu} \lambda_i g_i,$$

where $g_i \in \mathbf{C}\{z_0, \dots, z_n\}/(f_{z_0}, \dots, f_{z_n})$ are generators of the Jacobi ring, which are monomials. (Since f is quasi-homogeneous, g_i can be taken as monomials.) Then,

$$\begin{aligned} & \{\lambda = (\lambda_1, \dots, \lambda_\mu) \in U \subset \mathbf{C}^\mu ; \mu(f, 0) = \mu(f_\lambda, 0)\} \\ &= \{\lambda = (\lambda_1, \dots, \lambda_\mu) \in U \subset \mathbf{C}^\mu ; \lambda_i = 0 \text{ for all } i \text{ satisfying that } \deg_\alpha(g_i) < 1\}, \end{aligned}$$

where $\deg_\alpha(z_0^{i_0} \dots z_n^{i_n}) := \alpha_0 i_0 + \dots + \alpha_n i_n$. $\square$

From 1.7 (1) and 2.1, it follows that

$$\mu(f, 0) = \mu(f_\lambda, 0) \iff P_g(f, 0) = P_g(f_\lambda, 0)$$

holds true if f_λ is also nondegenerate. Though f_λ is not always nondegenerate, the following theorem is useful as well.

Theorem 2.2 (Reid [R1]-Thm. 4.1, [R2]-Thm. 4.6). *Let $(X = \{f = 0\}, 0) \subset (\mathbf{C}^{n+1}, 0)$ be a hypersurface singularity and let $\alpha = (\alpha_0, \dots, \alpha_n) = (p_0/p, \dots, p_n/p) \in \mathbf{Q}_{>0}^{n+1}$ such that $(p_0, \dots, p_n) \in \mathbf{N}^{n+1}$ is a primitive vector. Then,*

$$(X, 0) : \text{canonical} \implies \deg_\alpha(z_0 \cdots z_n) > \deg_\alpha(f),$$

where $\deg_\alpha(f) := \min\{\deg_\alpha(z^\nu) ; z^\nu \in f\}$. $\square$

A hypersurface singularity is canonical if and only if it is rational, and so it follows that

$$\mu(f, 0) = \mu(f_\lambda, 0) \implies P_g(f, 0) = P_g(f_\lambda, 0)$$

holds always true from 2.2. Hence we should show the converse proposition.

Remark 2.3. Let (X, x) be a n -dimensional normal Gorenstein singularity and

$$(\tilde{\tilde{X}}, \tilde{E}) \xrightarrow{\pi'} (\tilde{X}, E) \xrightarrow{\pi} (X, x),$$

where $\tilde{\tilde{X}}$ has at most rational singularities and $\pi'' := \pi \circ \pi'$ is a resolution. Let $E = \bigcup_{i \in I} E_i$ be the decomposition of the exceptional set E into irreducible components, and write $K_{\tilde{X}} = \pi^* K_X + \sum_{i \in I} m_i E_i$. Then, it follows that $\pi'_*(\mathcal{O}(K_{\tilde{\tilde{X}}})) = \mathcal{O}(K_{\tilde{X}})$ since $\tilde{\tilde{X}}$ has at most rational singularities ([KKMS]-p. 50), and so

$$\begin{aligned} P_g(X, x) &:= \dim_{\mathbf{C}}(R^{n-1} \pi''_* \mathcal{O}_{\tilde{\tilde{X}}})_x \\ &= \dim_{\mathbf{C}} \Gamma(X - \{x\}, \mathcal{O}(K_X)) / L^2(X - \{x\}) \\ &= \dim_{\mathbf{C}} \Gamma(\tilde{\tilde{X}} - \tilde{E}, \mathcal{O}(K_{\tilde{\tilde{X}}})) / \Gamma(\tilde{\tilde{X}}, \mathcal{O}(K_{\tilde{\tilde{X}}})) \\ &= \dim_{\mathbf{C}} \Gamma(\tilde{\tilde{X}}, \mathcal{O}(\pi''^* K_X)) / \Gamma(\tilde{\tilde{X}}, \mathcal{O}(K_{\tilde{\tilde{X}}})) \\ &= \dim_{\mathbf{C}} \Gamma(\tilde{X}, \mathcal{O}(\pi^* K_X)) / \Gamma(\tilde{X}, \pi'_*(\mathcal{O}(K_{\tilde{\tilde{X}}})) \\ &= \dim_{\mathbf{C}} \Gamma(\tilde{X}, \mathcal{O}(\pi^* K_X)) / \Gamma(\tilde{X}, \mathcal{O}(K_{\tilde{X}})). \end{aligned}$$

Therefore,

$$\exists i \in I ; m_i < 0 \iff \Gamma(\tilde{X}, \mathcal{O}(\pi^* K_X)) \supsetneq \Gamma(\tilde{X}, \mathcal{O}(K_{\tilde{X}})) \iff P_g(X, x) > 0,$$

namely,

$$\forall i \in I, m_i \geq 0 \iff P_g(X, x) = 0.$$

Theorem 2.4 (Tomari-Watanabe [TW]-Thm. 5.6). *Let $(X = \{f = 0\}, x)$ be a n -dimensional hypersurface isolated singularity and*

$$f = f_0 + f_1 + f_2 + \cdots,$$

$$f_i : \text{quasi-homogeneous polynomial of weight } \alpha = (\alpha_0, \cdots, \alpha_n);$$

$$1 = \deg_\alpha(f_0) < \deg_\alpha(f_1) < \deg_\alpha(f_2) < \cdots,$$

and

$$\begin{array}{ccc} V & \xrightarrow{\Pi : \alpha\text{-blow-up}} & \mathbf{C}^{n+1} \\ \cup & & \cup \\ \tilde{X} = \overline{\Pi^{-1}(X) - \Pi^{-1}(x)} & \xrightarrow{\pi = \Pi|_{\tilde{X}}} & X \\ \cup & & \cup \\ E = \pi^{-1}(x) & \longrightarrow & x. \end{array}$$

Assume that f_0 is irreducible and both $X - \{x\}$ and $\{f_0 = 0\} - \{x\}$ have at most rational singularities around x . Then $\tilde{X}$ has at most rational singularities. $\square$

Thus we expect a partial resolution π in 2.3 is given by a weighted blow-up.

Remark 2.5 (Ishii [I2]-Prop. 1.3, 1.6). Under the notation in 2.4, $\Pi^*X \subset V$ and $K_{\mathbf{C}^4}$ are principal divisors, hence,

$$\begin{aligned} \Pi^*X &= \tilde{X} + pF, \\ K_V &= \Pi^*K_{\mathbf{C}^4} + (p_1 + p_2 + p_3 + p_4 - 1)F, \\ K_X &= (K_{\mathbf{C}^4} + X)|_X, \\ \text{thus, } K_{\tilde{X}} &= (K_V + \tilde{X})|_{\tilde{X}} \\ &= (\Pi^*(K_{\mathbf{C}^4} + X) + (p_1 + p_2 + p_3 + p_4 - 1 - p)F)|_{\tilde{X}} \\ &= \pi^*K_X + (p_1 + p_2 + p_3 + p_4 - 1 - p) \sum_i k_i E_i, \end{aligned}$$

where $\alpha = (\alpha_1, \alpha_2, \alpha_3, \alpha_4) = (p_1/p, p_2/p, p_3/p, p_4/p)$ with $\mathbf{gcd}(p_1, p_2, p_3, p_4) = 1$,

$$F = \Pi^{-1}(0), \quad F|_{\tilde{X}} = E = \sum_i k_i E_i, \quad k_i > 0.$$

Thus, $p < p_1 + \cdots + p_4$ if and only if $R^2\pi_*\mathcal{O}_{\tilde{X}} = 0$.

Problem 2.6. Let f be a nondegenerate quasi-homogeneous polynomial which defines a simple $K3$ singularity at 0, and $(X_\lambda = \{f_\lambda = 0\}, 0)$ a versal deformation of $(X = \{f = 0\}, 0)$ such that $\mu(X_\lambda, 0) < \mu(X, 0)$. Then find a weight $\alpha' = (\alpha'_1, \dots, \alpha'_4) = (p'_1/p', \dots, p'_4/p')$ of f_λ with $1 < \alpha'_1 + \dots + \alpha'_4$ such that:

$$f_\lambda = f_0 + f_1 + f_2 + \dots,$$

$$f_i : \text{quasi-homogeneous polynomial of weight } \alpha',$$

$$1 = \deg_{\alpha'}(f_0) < \deg_{\alpha'}(f_1) < \deg_{\alpha'}(f_2) < \dots,$$

$$f_0 \text{ is irreducible,}$$

and $\{f_0 = 0\} - \{0\}$ has at most rational singularities around 0.

If there exists such weight α' then $P_g(X_\lambda, 0) = 0$ by 2.4 and 2.5.

We show there exists such weight α' as in 2.6 for $f = x^2 + y^3 + \dots$, of No.10-14, 46-51 and 83 in Table 2.2 of [Yo] to obtain our main result as follows:

Theorem 2.7. *Let $(X = \{f = 0\}, 0)$ be a hypersurface simple $K3$ singularity defined by a nondegenerate quasi-homogeneous polynomial $f = x^2 + y^3 + \dots$, which is one of No.10-14, 46-51 and 83 in Table 2.2 of [Yo]:*

$$(No. 10) \quad f = x^2 + y^3 + z^{12} + w^{12},$$

$$(No. 11) \quad f = x^2 + y^3 + z^{10} + w^{15},$$

$$(No. 12) \quad f = x^2 + y^3 + z^9 + w^{18},$$

$$(No. 13) \quad f = x^2 + y^3 + z^8 + w^{24},$$

$$(No. 14) \quad f = x^2 + y^3 + z^7 + w^{42},$$

$$(No. 46) \quad f = x^2 + y^3 + z^{11} + zw^{12},$$

$$(No. 47) \quad f = x^2 + y^3 + yz^7 + z^9w^2 + w^{14},$$

$$(No. 48) \quad f = x^2 + y^3 + z^9w + w^{16},$$

$$(No. 49) \quad f = x^2 + y^3 + z^8w + w^{21},$$

$$(No. 50) \quad f = x^2 + y^3 + yz^5 + z^7w^2 + w^{30},$$

$$(No. 51) \quad f = x^2 + y^3 + z^7w + w^{36},$$

$$(No. 83) \quad f = x^2 + y^3 + yw^9 + z^{10}w + z^2w^{11},$$

and let $\mathfrak{X}$ be the versal deformation of $(X, 0)$. Then,

$$\begin{aligned} & \{\lambda \in U \subset \mathbf{C}^N ; \mu(X, 0) = \mu(X_\lambda, 0), (X_\lambda, 0) \in \mathfrak{X}\} \\ &= \{\lambda \in U \subset \mathbf{C}^N ; 1 = \delta_m(X, 0) = \delta_m(X_\lambda, 0) \text{ for all } m \geq 1, (X_\lambda, 0) \in \mathfrak{X}\}. \end{aligned}$$

3. PROOF OF THEOREM 2.7

We prepare the following Lemma prior to the proof of Theorem 2.7.

Lemma 3.1. *Let f be a nondegenerate quasi-homogeneous polynomial listed by Yonemura [Yo] which defines a simple K3 singularity, and f_λ a versal deformation of f such that $\mu(f_\lambda, 0) < \mu(f, 0)$. Under this original local coordinate system, if $\alpha' = (\alpha'_1, \dots, \alpha'_4) \in \mathbf{Q}_{>0}^4$ satisfies $\deg_{\alpha'}(f_\lambda) = 1$ then $1 < \alpha'_1 + \dots + \alpha'_4$.*

Proof. For any $i=1,2,3,4$, one of the following is satisfied:

- (a) $p_i | p$,
- (b) $p_i | (p - p_j)$ for some $j \neq i$,

namely,

- (a) $z_i^I \in f \quad (\exists I \geq 2)$,
- (b) $z_i^I z_j \in f \quad (\exists I \geq 2)$ for some $j \neq i$.

By assumptions, $p\alpha_1 + q\alpha_2 + r\alpha_3 + s\alpha_4 < 1$ for some $z_1^p z_2^q z_3^r z_4^s \in f_\lambda$,

and $p\alpha'_1 + q\alpha'_2 + r\alpha'_3 + s\alpha'_4 \geq 1$ for all $z_1^p z_2^q z_3^r z_4^s \in f_\lambda$,

so there exists $i \in \{1, 2, 3, 4\}$ such that $\alpha_i < \alpha'_i \quad \dots \circledast$.

So the defining polynomials f can be classified as below.

Case 1. $z_i^I \in f \quad (\exists I \geq 2)$.

Then $I\alpha_i = 1 \leq I\alpha'_i$, so we have $\alpha_i \leq \alpha'_i$.

(1-I) $z_i^I, z_j^J \in f \quad (\exists I, \exists J \geq 2)$.

Since $J\alpha_j = 1 \leq J\alpha'_j$, we have $\alpha_j \leq \alpha'_j$.

(1-I-i) $z_i^I, z_j^J, z_k^K \in f \quad (\exists I, \exists J, \exists K \geq 2)$.

From $K\alpha_k = 1 \leq K\alpha'_k$, we have $\alpha_k \leq \alpha'_k$.

(1-I-i-a) (No.1-14) $z_i^I, z_j^J, z_k^K, z_l^L \in f \quad (\exists I, \exists J, \exists K, \exists L \geq 2)$.

From $L\alpha_l = 1 \leq L\alpha'_l$, we have $\alpha_l \leq \alpha'_l$, so $\alpha_1 + \dots + \alpha_4 < \alpha'_1 + \dots + \alpha'_4$ by $\circledast$.

(1-I-i-b) (No.15-51) $z_i^I, z_j^J, z_k^K, z_k z_l^L \in f \quad (\exists I, \exists J, \exists K, \exists L \geq 2)$.

From $\alpha_k + L\alpha_l = 1 \leq \alpha'_k + L\alpha'_l$, we have $\alpha_k + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_k + \alpha'_l$.

If $\alpha_l \leq \alpha'_l$ then $\alpha_1 + \dots + \alpha_4 < \alpha'_1 + \dots + \alpha'_4$ by $\circledast$.

If $\alpha_l > \alpha'_l$ then

$\alpha_i + \alpha_j + \alpha_k + \alpha_l < \alpha_i + \alpha_j + \alpha_k + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_i + \alpha'_j + \alpha'_k + \alpha'_l$.

(1-I-ii) $z_i^I, z_j^J, z_k^K z_l \in f \quad (\exists I, \exists J, \exists K \geq 2)$.

$\alpha_k \leq \alpha'_k$ or $\alpha_l \leq \alpha'_l$, and $\alpha_k + \alpha_l + (K-1)(\alpha_k - \alpha'_k) \leq \alpha'_k + \alpha'_l$.

(1-I-ii-a) (No.78) $z_i^I, z_j^J, z_k^K z_l, z_k z_l^L \in f \quad (\exists I, \exists J, \exists K, \exists L \geq 2).$

$$\alpha_k + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_k + \alpha'_l.$$

If $\alpha_k \leq \alpha'_k$ and $\alpha_l \leq \alpha'_l$ then $\alpha_1 + \dots + \alpha_4 < \alpha'_1 + \dots + \alpha'_4$ by $\circledast$.

If $\alpha_k \leq \alpha'_k$ and $\alpha_l > \alpha'_l$ then

$$\alpha_i + \alpha_j + \alpha_k + \alpha_l < \alpha_i + \alpha_j + \alpha_k + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_i + \alpha'_j + \alpha'_k + \alpha'_l.$$

If $\alpha_k > \alpha'_k$ and $\alpha_l \leq \alpha'_l$ then

$$\alpha_i + \alpha_j + \alpha_k + \alpha_l < \alpha_i + \alpha_j + \alpha_k + \alpha_l + (K-1)(\alpha_k - \alpha'_k) \leq \alpha'_i + \alpha'_j + \alpha'_k + \alpha'_l.$$

(1-I-ii-b) (No.52, 54-74, 76, 77, 79-83)

$$z_i^I, z_j^J, z_k^K z_l, z_j z_l^L \in f \quad (\exists I, \exists J, \exists K, \exists L \geq 2).$$

$\alpha_j + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_j + \alpha'_l$. The rest is similar to 1-I-ii-a.

(1-I-iii) $z_i^I, z_j^J, z_j z_k^K \in f \quad (\exists I, \exists J, \exists K \geq 2).$

Similarly, $\alpha_j + \alpha_k + (K-1)(\alpha_k - \alpha'_k) \leq \alpha'_j + \alpha'_k$.

(1-I-iii-a) (No.66, 67, 72, 75, 81, 82)

$$z_i^I, z_j^J, z_j z_k^K, z_j z_l^L \in f \quad (\exists I, \exists J, \exists K, \exists L \geq 2).$$

Since there exists $z_k^r z_l^s \in f$, we have $\alpha_k \leq \alpha'_k$ or $\alpha_l \leq \alpha'_l$.

Moreover $\alpha_j + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_j + \alpha'_l$ by $z_j z_l^L \in f$.

So we have $\alpha_1 + \dots + \alpha_4 < \alpha'_1 + \dots + \alpha'_4$ similarly as in 1-I-ii-a.

(1-I-iii-b) (No.53, 57, 58, 62-64, 66, 67, 69-72)

$$z_i^I, z_j^J, z_j z_k^K, z_i z_l^L \in f \quad (\exists I, \exists J, \exists K, \exists L \geq 2).$$

$$\alpha_i + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_i + \alpha'_l.$$

If $\alpha_k \leq \alpha'_k$ or $\alpha_l \leq \alpha'_l$ then the rest is similar to 1-I-ii-a.

If $\alpha_k > \alpha'_k$ and $\alpha_l > \alpha'_l$ then

$$\begin{aligned} \alpha_i + \alpha_j + \alpha_k + \alpha_l &< \alpha_i + \alpha_j + \alpha_k + \alpha_l + (K-1)(\alpha_k - \alpha'_k) + (L-1)(\alpha_l - \alpha'_l) \\ &\leq \alpha'_i + \alpha'_j + \alpha'_k + \alpha'_l. \end{aligned}$$

(1-II) $z_i^I, z_j^J z_k \in f \quad (\exists I, \exists J \geq 2).$

$$\alpha_j \leq \alpha'_j \text{ or } \alpha_k \leq \alpha'_k, \text{ and } \alpha_j + \alpha_k + (J-1)(\alpha_j - \alpha'_j) \leq \alpha'_j + \alpha'_k.$$

(1-II-i) $z_i^I, z_j^J z_k, z_k^K z_l \in f \quad (\exists I, \exists J, \exists K \geq 2).$

$$\alpha_k \leq \alpha'_k \text{ or } \alpha_l \leq \alpha'_l, \text{ and } \alpha_k + \alpha_l + (K-1)(\alpha_k - \alpha'_k) \leq \alpha'_k + \alpha'_l.$$

(1-II-i-a) (No.88, 90-93) $z_i^I, z_j^J z_k, z_k^K z_l \in f \quad (\exists I, \exists J, \exists K, \exists L \geq 2).$

Since there exists $z_j^q z_l^s \in f$, we have $\alpha_j \leq \alpha'_j$ or $\alpha_l \leq \alpha'_l$.

Moreover $\alpha_k + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_k + \alpha'_l$ by $z_k z_l^L \in f$.

If $\alpha_j \leq \alpha'_j$, $\alpha_k \leq \alpha'_k$ and $\alpha_l \leq \alpha'_l$, then the assertion holds true.

If $\alpha_j \leq \alpha'_j$, $\alpha_k \leq \alpha'_k$ and $\alpha_l > \alpha'_l$, then

$$\alpha_i + \alpha_j + \alpha_k + \alpha_l < \alpha_i + \alpha_j + \alpha_k + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_i + \alpha'_j + \alpha'_k + \alpha'_l.$$

The assertion holds true similarly for both the case of $\alpha_j \leq \alpha'_j$, $\alpha_k > \alpha'_k$,

$\alpha_l \leq \alpha'_l$, and the case of $\alpha_j > \alpha'_j$, $\alpha_k \leq \alpha'_k$, $\alpha_l \leq \alpha'_l$.

(1-II-i-b) (No.86-88, 92) $z_i^I, z_j^J z_k, z_k^K z_l, z_j z_l^L \in f \quad (\exists I, \exists J, \exists K, \exists L \geq 2).$

$$\alpha_j \leq \alpha'_j \text{ or } \alpha_l \leq \alpha'_l, \text{ and } \alpha_j + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_j + \alpha'_l.$$

The rest is similar to 1-II-i-a.

(1-II-i-c) (No.84-89) $z_i^I, z_j^J z_k, z_k^K z_l, z_i z_l^L \in f \quad (\exists I, \exists J, \exists K, \exists L \geq 2).$

$$\alpha_i + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_i + \alpha'_l.$$

When $\alpha_k \leq \alpha'_k$, the rest is similar to 1-I-iii-b.

If $\alpha_k > \alpha'_k$ then $\alpha_j \leq \alpha'_j$ and $\alpha_l \leq \alpha'_l$, so

$$\alpha_i + \alpha_j + \alpha_k + \alpha_l < \alpha_i + \alpha_j + \alpha_k + \alpha_l + (K-1)(\alpha_k - \alpha'_k) \leq \alpha'_i + \alpha'_j + \alpha'_k + \alpha'_l.$$

(1-II-ii) $z_i^I, z_j^J z_k, z_j z_k^K \in f \quad (\exists I, \exists J, \exists K \geq 2).$

$$\alpha_j + \alpha_k + (K-1)(\alpha_k - \alpha'_k) \leq \alpha'_j + \alpha'_k.$$

(1-II-ii-a) (No.89) $z_i^I, z_j^J z_k, z_j z_k^K, z_i z_l^L \in f \quad (\exists I, \exists J, \exists K, \exists L \geq 2).$

$$\alpha_i + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_i + \alpha'_l.$$

For the case $\alpha_l \leq \alpha'_l$, the assertion holds true.

If $\alpha_l > \alpha'_l$ then

$$\begin{aligned} & \alpha_i + \alpha_j + \alpha_k + \alpha_l < \alpha_i + \alpha_j + \alpha_k + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \\ & < \begin{cases} \alpha_i + \alpha_j + \alpha_k + \alpha_l + (J-1)(\alpha_j - \alpha'_j) + (L-1)(\alpha_l - \alpha'_l) & (\text{if } \alpha_j > \alpha'_j) \\ \alpha_i + \alpha_j + \alpha_k + \alpha_l + (K-1)(\alpha_k - \alpha'_k) + (L-1)(\alpha_l - \alpha'_l) & (\text{if } \alpha_k > \alpha'_k) \end{cases} \\ & \leq \alpha'_i + \alpha'_j + \alpha'_k + \alpha'_l. \end{aligned}$$

(1-II-iii) $z_i^I, z_j^J z_k, z_i z_k^K \in f \quad (\exists I, \exists J, \exists K \geq 2).$

$$\alpha_i + \alpha_k + (K-1)(\alpha_k - \alpha'_k) \leq \alpha'_i + \alpha'_k.$$

(1-II-iii-a) (No.89) $z_i^I, z_j^J z_k, z_i z_k^K, z_k z_l^L \in f \quad (\exists I, \exists J, \exists K, \exists L \geq 2).$

$$\alpha_k \leq \alpha'_k \text{ or } \alpha_l \leq \alpha'_l, \text{ and } \alpha_k + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_k + \alpha'_l.$$

Moreover $\alpha_j \leq \alpha'_j$ or $\alpha_l \leq \alpha'_l$ since there exists $z_j^q z_l^s \in f$.

So $\alpha_1 + \dots + \alpha_4 < \alpha'_1 + \dots + \alpha'_4$ similarly as 1-II-i-a.

(1-II-iii-b) (No.85, 87, 89) $z_i^I, z_j^J z_k, z_i z_k^K, z_i z_l^L \in f \quad (\exists I, \exists J, \exists K, \exists L \geq 2).$

Since there exists $z_k^r z_l^s \in f$, we have $\alpha_k \leq \alpha'_k$ or $\alpha_l \leq \alpha'_l$.

Moreover $\alpha_i + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_i + \alpha'_l$ by $z_i z_l^L \in f$.

When $\alpha_k \leq \alpha'_k$, the rest is similar to 1-I-iii-b.

If $\alpha_k > \alpha'_k$ then $\alpha_j \leq \alpha'_j$ and $\alpha_l \leq \alpha'_l$, so

$$\alpha_i + \alpha_j + \alpha_k + \alpha_l < \alpha_i + \alpha_j + \alpha_k + \alpha_l + (K-1)(\alpha_k - \alpha'_k) \leq \alpha'_i + \alpha'_j + \alpha'_k + \alpha'_l.$$

Case 2. $z_i^I z_j \in f$ ($\exists I \geq 2$).

$$\alpha_i \leq \alpha'_i \text{ or } \alpha_j \leq \alpha'_j, \text{ and } \alpha_i + \alpha_j + (I-1)(\alpha_i - \alpha'_i) \leq \alpha'_i + \alpha'_j.$$

(2-I) $z_i^I z_j, z_j^J z_k \in f$ ($\exists I, \exists J \geq 2$).

$$\alpha_j \leq \alpha'_j \text{ or } \alpha_k \leq \alpha'_k, \text{ and } \alpha_j + \alpha_k + (J-1)(\alpha_j - \alpha'_j) \leq \alpha'_j + \alpha'_k.$$

(2-I-i) $z_i^I z_j, z_j^J z_k, z_k^K z_l \in f$ ($\exists I, \exists J, \exists K \geq 2$).

$$\alpha_k \leq \alpha'_k \text{ or } \alpha_l \leq \alpha'_l, \text{ and } \alpha_k + \alpha_l + (K-1)(\alpha_k - \alpha'_k) \leq \alpha'_k + \alpha'_l.$$

(2-I-i-a) (No.94, 95) $z_i^I z_j, z_j^J z_k, z_k^K z_l, z_k z_l^L \in f$ ($\exists I, \exists J, \exists K, \exists L \geq 2$).

$$\alpha_k + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_k + \alpha'_l.$$

Since there exists $z_j^q z_l^s \in f$, we have $\alpha_j \leq \alpha'_j$ or $\alpha_l \leq \alpha'_l$.

If $\alpha_i \leq \alpha'_i$ then the rest is similar to 1-II-i-a.

If $\alpha_i > \alpha'_i$ then $\alpha_j \leq \alpha'_j$ and so

$$\begin{aligned} \alpha_i + \alpha_j + \alpha_k + \alpha_l &< \alpha_i + \alpha_j + \alpha_k + \alpha_l + (I-1)(\alpha_i - \alpha'_i) \\ &< \begin{cases} \alpha_i + \alpha_j + \alpha_k + \alpha_l + (I-1)(\alpha_i - \alpha'_i) + (K-1)(\alpha_k - \alpha'_k) & (\text{if } \alpha_k > \alpha'_k) \\ \alpha_i + \alpha_j + \alpha_k + \alpha_l + (I-1)(\alpha_i - \alpha'_i) + (L-1)(\alpha_l - \alpha'_l) & (\text{if } \alpha_l > \alpha'_l) \end{cases} \\ &\leq \alpha'_i + \alpha'_j + \alpha'_k + \alpha'_l. \end{aligned}$$

(2-I-i-b) (No.94, 95) $z_i^I z_j, z_j^J z_k, z_k^K z_l, z_j z_l^L \in f$ ($\exists I, \exists J, \exists K, \exists L \geq 2$).

$$\alpha_j \leq \alpha'_j \text{ or } \alpha_l \leq \alpha'_l, \text{ and } \alpha_j + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_j + \alpha'_l.$$

Since there exists $z_i^p z_l^s \in f$, we have $\alpha_i \leq \alpha'_i$ or $\alpha_l \leq \alpha'_l$.

If $\alpha_i \leq \alpha'_i$ then the rest is similar to 1-II-i-a.

If $\alpha_i > \alpha'_i$ then $\alpha_j \leq \alpha'_j$ and $\alpha_l \leq \alpha'_l$, and so

$$\begin{aligned} \alpha_i + \alpha_j + \alpha_k + \alpha_l &< \alpha_i + \alpha_j + \alpha_k + \alpha_l + (I-1)(\alpha_i - \alpha'_i) + (K-1)(\alpha_k - \alpha'_k) \\ &\leq \alpha'_i + \alpha'_j + \alpha'_k + \alpha'_l \quad \text{for the case of } \alpha_k > \alpha'_k. \end{aligned}$$

(2-I-i-c) (No.94, 95) $z_i^I z_j, z_j^J z_k, z_k^K z_l, z_i z_l^L \in f$ ($\exists I, \exists J, \exists K, \exists L \geq 2$).

$$\alpha_i \leq \alpha'_i \text{ or } \alpha_l \leq \alpha'_l, \text{ and } \alpha_i + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_i + \alpha'_l.$$

If $\alpha_i \leq \alpha'_i$ and $\alpha_j \leq \alpha'_j$ then the assertion holds true.

If $\alpha_i \leq \alpha'_i$ and $\alpha_j > \alpha'_j$ then $\alpha_k \leq \alpha'_k$, so the rest is similar to 1-I-iii-b.

If $\alpha_i > \alpha'_i$ and $\alpha_j \leq \alpha'_j$ then $\alpha_l \leq \alpha'_l$, so the rest is similar to 1-I-iii-b.

(2-II) $z_i^I z_j, z_i z_j^J \in f$ ($\exists I, \exists J \geq 2$).

$$\alpha_i + \alpha_j + (J-1)(\alpha_j - \alpha'_j) \leq \alpha'_i + \alpha'_j.$$

(2-II-i) $z_i^I z_j, z_i z_j^J, z_k^K z_l \in f$ ($\exists I, \exists J, \exists K \geq 2$).

$$\begin{aligned}
& \alpha_k \leq \alpha'_k \text{ or } \alpha_l \leq \alpha'_l, \text{ and } \alpha_k + \alpha_l + (K-1)(\alpha_k - \alpha'_k) \leq \alpha'_k + \alpha'_l. \\
\textbf{(2-II-i-a)} \quad & z_i^I z_j, z_i z_j^J, z_k^K z_l, z_k z_l^L \in f \quad (\exists I, \exists J, \exists K, \exists L \geq 2). \\
& \alpha_k + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_k + \alpha'_l. \\
& \alpha_i + \alpha_j + \alpha_k + \alpha_l \\
& < \begin{cases} \alpha_i + \cdots + \alpha_l + (I-1)(\alpha_i - \alpha'_i) & (\text{if } \alpha_i > \alpha'_i) \\ \dots\dots\dots & \dots\dots \\ \alpha_i + \cdots + \alpha_l + (L-1)(\alpha_l - \alpha'_l) & (\text{if } \alpha_l > \alpha'_l) \end{cases} \\
& < \begin{cases} \alpha_i + \cdots + \alpha_l + (I-1)(\alpha_i - \alpha'_i) + (K-1)(\alpha_k - \alpha'_k) & (\alpha_i > \alpha'_i, \alpha_k > \alpha'_k) \\ \alpha_i + \cdots + \alpha_l + (I-1)(\alpha_i - \alpha'_i) + (L-1)(\alpha_l - \alpha'_l) & (\alpha_i > \alpha'_i, \alpha_l > \alpha'_l) \\ \alpha_i + \cdots + \alpha_l + (J-1)(\alpha_j - \alpha'_j) + (K-1)(\alpha_k - \alpha'_k) & (\alpha_j > \alpha'_j, \alpha_k > \alpha'_k) \\ \alpha_i + \cdots + \alpha_l + (J-1)(\alpha_j - \alpha'_j) + (L-1)(\alpha_l - \alpha'_l) & (\alpha_j > \alpha'_j, \alpha_l > \alpha'_l) \end{cases} \\
& \leq \alpha'_i + \alpha'_j + \alpha'_k + \alpha'_l. \\
\textbf{(2-II-ii)} \quad & z_i^I z_j, z_i z_j^J, z_j z_k^K \in f \quad (\exists I, \exists J, \exists K \geq 2). \\
& \alpha_j \leq \alpha'_j \text{ or } \alpha_k \leq \alpha'_k, \text{ and } \alpha_j + \alpha_k + (K-1)(\alpha_k - \alpha'_k) \leq \alpha'_j + \alpha'_k. \\
\textbf{(2-II-ii-a)} \quad & z_i^I z_j, z_i z_j^J, z_j z_k^K, z_j z_l^L \in f \quad (\exists I, \exists J, \exists K, \exists L \geq 2). \\
& \alpha_j \leq \alpha'_j \text{ or } \alpha_l \leq \alpha'_l, \text{ and } \alpha_j + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_j + \alpha'_l. \\
\textbf{(2-II-ii-b)} \quad & \text{(No.94, 95)} \quad z_i^I z_j, z_i z_j^J, z_j z_k^K, z_i z_l^L \in f \quad (\exists I, \exists J, \exists K, \exists L \geq 2). \\
& \alpha_i \leq \alpha'_i \text{ or } \alpha_l \leq \alpha'_l, \text{ and } \alpha_i + \alpha_l + (L-1)(\alpha_l - \alpha'_l) \leq \alpha'_i + \alpha'_l. \\
& \text{Since there exists } z_i^p z_k^r \in f, \text{ we have } \alpha_i \leq \alpha'_i \text{ or } \alpha_k \leq \alpha'_k. \\
& \text{Since there exists } z_j^q z_l^s \in f, \text{ we have } \alpha_j \leq \alpha'_j \text{ or } \alpha_l \leq \alpha'_l. \\
& \text{If } \alpha_l \leq \alpha'_l \text{ then the rest is similar to } \textbf{1-II-i-a}. \\
& \text{If } \alpha_l > \alpha'_l \text{ then } \alpha_i \leq \alpha'_i \text{ and } \alpha_j \leq \alpha'_j, \text{ so the rest is similar to } \textbf{2-I-i-b}.
\end{aligned}$$

Q.E.D.

Lemma 3.2. Let $f = \alpha(x, y, z)x^2 + \beta(y, z)y^3 + \phi(z)y^2 + \varphi(z)y + \psi(z) \in \mathbf{C}[x, y, z]$ define an isolated singularity at the origin $0 \in \mathbf{C}^3$, which satisfies one of the following:

- (1) $\beta(y, z) = \beta_0 + \text{higher terms}$, $0 \neq \beta_0 \in \mathbf{C}$, $\text{ord}\phi = 1$, and $\beta_0 y^3 + \phi_0(z)y^2 + \varphi_0(z)y + \psi_0(z)$ has no triple factor,
- (1') $\beta \equiv 0$ or $\text{ord}\beta \geq 1$, and $\text{ord}\phi = 1$,
- (2) $\beta(y, z) = \beta_0 + \text{higher terms}$, $0 \neq \beta_0 \in \mathbf{C}$, $\text{ord}\phi \leq 3$ or $\text{ord}\psi \leq 5$, and $\beta_0 y^3 + \phi_0(z)y^2 + \varphi_0(z)y + \psi_0(z)$ has no triple factor,

where $\text{ord}\alpha = 0$, and $\phi_0, \varphi_0, \psi_0$ are the initial parts of ϕ, φ, ψ , respectively. Then $(f, 0)$ is rational.

Proof. We may assume that $\alpha(x, y, z) = 1, \beta_0 = 1$.

(1) If $\text{ord}\varphi \leq 1$ or $\text{ord}\psi \leq 2$ then the assertion holds true. So we may assume $\text{ord}\varphi \geq 2$ and $\text{ord}\psi \geq 3$. Let f_0 be the initial part of f with respect to the weight $\alpha = (1/2, p, q)$, where α satisfies the conditions

$$\frac{1}{3} \leq p = \frac{1-q}{2} < \frac{1}{2} < p+q \left(= \frac{1+q}{2} \right),$$

$$\deg_\alpha(\varphi(z)y), \deg_\alpha\psi(z) \geq 1, \quad \text{and} \quad \deg_\alpha(\varphi(z)y) \text{ or } \deg_\alpha\psi(z) = 1.$$

(There exists such weight α because

$$\frac{1-q}{2} \geq \frac{1}{3}, \quad \frac{1-q}{2} + Nq \geq 1 \quad \text{for } N \geq 2, \quad \frac{1}{3} \geq q \geq \frac{1}{2N-1}.)$$

If $f_0 - x^2 \in \mathbf{C}[y, z]$ has no double factor, then $(f_0, 0)$ is an isolated singularity, so $(f_0, 0)$ is rational. So $\{f_0 = 0\} \subset \mathbf{C}^3$ has only rational singularities around the origin $0 \in \mathbf{C}^3$. Therefore $(f, 0)$ is also rational from 2.4 and 2.5, because $\frac{1}{2} + p + q > 1$.

If $f_0 - x^2$ has a double factor, namely,

$$f = x^2 + (y + \gamma_1 z)^2(y + \gamma_2 z) + \text{higher terms}, \quad \gamma_1 \neq \gamma_2,$$

then taking the coordinate changes $Y := y + \gamma_1 z$ and $z' := (\gamma_2 - \gamma_1)z$, thus

$$f = x^2 + \beta'(Y, z')Y^3 + \phi'(z')Y^2 + \varphi'(z')Y + \psi'(z'),$$

for some $\beta' \in \mathbf{C}[Y, z']$, $\phi', \varphi', \psi' \in \mathbf{C}[z']$ with $\beta' = 1 + \text{higher terms}$, $\phi' = z' + \text{higher terms}$, $2 = \text{ord}\varphi < \text{ord}\varphi'$, $3 = \text{ord}\psi < \text{ord}\psi'$. Let φ'_0, ψ'_0 be the initial parts of φ', ψ' , respectively. We replace the weight $\alpha = (1/2, p, q)$ with $\alpha' = (1/2, p', q')$, which satisfies:

$$\frac{1}{3} = p < p' = \frac{1-q'}{2} < \frac{1}{2} < p' + q' \left(= \frac{1+q'}{2} \right),$$

$$\deg_{\alpha'}(\varphi'(z')Y), \deg_{\alpha'}\psi'(z') \geq 1, \quad \text{and} \quad \deg_{\alpha'}(\varphi'(z')Y) \text{ or } \deg_{\alpha'}\psi'(z') = 1.$$

(There exists such weight α' because

$$\frac{1-q'}{2} > \frac{1}{3}, \quad \frac{1-q'}{2} + Nq' \geq 1 \quad \text{for } N > 2, \quad \frac{1}{3} > q' \geq \frac{1}{2N-1}.)$$

If $Y^2 z' + Y\varphi'_0(z') + \psi'_0(z') = (Y + g(z'))^2 z'$ for some $g \in \mathbf{C}[z']$ ($\text{ord}(g) > 1$),

then taking the coordinate change $Y' := Y + g(z')$, thus

$$f = x^2 + \beta''(Y', z')Y'^3 + \phi''(z')Y'^2 + \varphi''(z')Y' + \psi''(z'),$$

for some $\beta'' \in \mathbf{C}[Y', z']$, $\phi'', \varphi'', \psi'' \in \mathbf{C}[z']$ with $\beta'' = 1 + \text{higher terms}$, $\phi'' =$

$z' + \text{higher terms}$, $\text{ord}\varphi' < \text{ord}\varphi''$, $\text{ord}\psi' < \text{ord}\psi''$. Let φ_0'' , ψ_0'' be the initial parts of φ'' , ψ'' , respectively. If

$$Y'^2 z' + Y' \varphi_0''(z') + \psi_0''(z') = (Y' + g'(z'))^2 z'$$

for some $g' \in \mathbf{C}[z']$ ($1 < \text{ord}(g) < \text{ord}(g')$), then taking the coordinate change

$Y'' := Y' + g'(z')$, $\dots$. If this procedure continues infinitely, then

$$\begin{aligned} \frac{1}{3} &= p < p' < \dots < p^{(n)} = \frac{1 - q^{(n)}}{2} < \dots < \frac{1}{2} < p^{(n)} + q^{(n)}, \\ \frac{1}{3} &= q > q' > \dots > q^{(n)} > \dots > 0, \end{aligned}$$

$$\dim_{\mathbf{C}} \mathbf{C}\{z'\} \left/ \left(\varphi^{(n)}, \frac{\partial \psi^{(n)}}{\partial z'} \right) \right. \leq \mu(f, 0) \xrightarrow{n \rightarrow +\infty} +\infty,$$

a contradiction. Therefore $(f, 0)$ is rational by 2.4 and 2.5.

(1') We may assume $\phi = z + \text{higher terms}$, $\text{ord}\varphi \geq 2$ and $\text{ord}\psi \geq 3$. Let f_0 be the initial part of f with respect to the weight $\alpha = (1/2, p, q)$, where α satisfies:

$$\frac{1}{3} \leq p = \frac{1 - q}{2} < \frac{1}{2} < p + q \left(= \frac{1 + q}{2} \right),$$

$$\deg_{\alpha}(\varphi(z)y), \deg_{\alpha}\psi(z) \geq 1, \quad \text{and} \quad \deg_{\alpha}(\varphi(z)y) \text{ or } \deg_{\alpha}\psi(z) = 1.$$

If $f_0 - x^2 \in \mathbf{C}[y, z]$ has no double factor, then $(f_0, 0)$ is rational similarly as in (1).

If $f_0 - x^2$ has a double factor, namely,

$$y^2 z + y\varphi_0(z) + \psi_0(z) = (y + g(z))^2 z \quad \text{for some } g \in \mathbf{C}[z] \quad (\text{ord}(g) \geq 1),$$

(where φ_0 , ψ_0 are the initial parts of φ , ψ , respectively,) then taking the coordinate change $Y := y + g(z)$, thus

$$f = \begin{cases} x^2 + \phi(z)Y^2 + \varphi'(z)Y + \psi'(z), & (\beta \equiv 0) \\ x^2 + \beta'(Y, z)Y^3 + \phi'(z)Y^2 + \varphi'(z)Y + \psi'(z), & (\text{ord}\beta \geq 1) \end{cases}$$

for some $\beta' \in \mathbf{C}[Y, z]$, ϕ' , φ' , $\psi' \in \mathbf{C}[z]$ with $1 \leq \text{ord}\beta'$, $\phi' = z + \text{higher terms}$, $2 \leq \text{ord}\varphi < \text{ord}\varphi'$, $3 \leq \text{ord}\psi < \text{ord}\psi'$. We replace the weight $\alpha = (1/2, p, q)$ with $\alpha' = (1/2, p', q')$, which satisfies the conditions

$$\frac{1}{3} \leq p < p' = \frac{1 - q'}{2} < \frac{1}{2} < p' + q' \left(= \frac{1 + q'}{2} \right),$$

$$\deg_{\alpha'}(\varphi'(z)Y), \deg_{\alpha'}\psi'(z) \geq 1, \quad \text{and} \quad \deg_{\alpha'}(\varphi'(z)Y) \text{ or } \deg_{\alpha'}\psi'(z) = 1.$$

If this procedure continues infinitely, then $\mu(f, 0) \xrightarrow{n \rightarrow +\infty} +\infty$, a contradiction.

(2) Let f_0 be the initial part of f with respect to the weight $\alpha = (1/2, 1/3, q)$.

If $f_0 - x^2 \in \mathbf{C}[y, z]$ has no double factor, then $(f_0, 0)$ is an isolated singularity, and so

$(f_0, 0)$ is rational for the case of $\text{ord}\varphi \leq 3$ or $\text{ord}\psi \leq 5$. Therefore $(f, 0)$ is also rational from 2.4 and 2.5, because $\frac{1}{2} + \frac{1}{3} + q > 1$ for $q > \frac{1}{6}$.

So consider the case of $f = x^2 + (y + \gamma_1 z)^2(y + \gamma_2 z) + \text{higher terms}$, $\gamma_1 \neq \gamma_2$.

Then the situation is similar to (1).

Q.E.D.

Example 3.3. (See Theorem 2.7.)

$$f = x^2 + y^3 + z^9 + w^{18}, \quad (\text{No.12})$$

$$f_\lambda = x^2 + y^3 + \left(-\frac{27}{4}c_{30}^2(z + \gamma_1 w)^4(z - 2\gamma_1 w)^2 + \varphi(z, w)\right)y \\ + (z + \gamma_1 w)^6(z - 2\gamma_1 w)^3 + \psi(z, w),$$

where $-\frac{27}{4}c_{30}^3 = 1$, $\text{ord}\varphi \geq 7$, $\text{ord}\psi \geq 10$, $\psi(z, w) \ni w^{18}$ (Case 10-b).

Taking the coordinate change $Y := y + \frac{3}{2}c_{30}(z + \gamma_1 w)^2(z - 2\gamma_1 w)$, we have

$$f_\lambda = x^2 + Y^3 - \frac{9}{2}c_{30}(z + \gamma_1 w)^2(z - 2\gamma_1 w)Y^2 + \varphi(z, w)Y \\ - \frac{3}{2}c_{30}(z + \gamma_1 w)^2(z - 2\gamma_1 w)\varphi(z, w) + \psi(z, w),$$

and then $z' := z + \gamma_1 w$,

$$f_\lambda = x^2 + Y^3 - \frac{9}{2}c_{30}(z'^3 - 3\gamma_1 z'^2 w)Y^2 + \varphi'(z', w)Y + \psi'(z', w)$$

for some $\varphi', \psi' \in \mathbf{C}[z', w]$, $\text{ord}\varphi' \geq 7$, $\text{ord}\psi' \geq 10$.

Let $\alpha' := (1/2, 1/3, 1/8, 1/12)$, $\alpha'' := (1/2, 17/50, 3/25, 2/25)$, $c'_{21} := -3\gamma_1 c_{30}$, and

$$\varphi' = -3(3c'_{21}z'^2 w + c'_{04}w^4)c'_{04}w^4 + \Phi', \quad \Phi'_0 = c'_{17}z'w^7, \\ \psi' = \left(-\frac{9}{2}c'_{21}z'^2 w - 2c'_{04}w^4\right)(c'_{04}w^4)^2 + \Psi', \quad \Psi'_0 = c'_{55}z'^5 w^5 + c'_{111}z'w^{11},$$

where Φ'_0, Ψ'_0 are the initial parts of Φ', Ψ' with respect to the weight α' . Then

$$f_\lambda = x^2 + Y^3 - \frac{9}{2}(c_{30}z'^3 + c'_{21}z'^2 w)Y^2 + (-3(3c'_{21}z'^2 w + c'_{04}w^4)c'_{04}w^4 + \Phi')(z', w)Y \\ + \left(\left(-\frac{9}{2}c'_{21}z'^2 w - 2c'_{04}w^4\right)(c'_{04}w^4)^2 + \Psi'\right)(z', w) \\ = x^2 + (Y + c'_{04}w^4)^2(Y - \frac{9}{2}c'_{21}z'^2 w - 2c'_{04}w^4) \\ - \frac{9}{2}c_{30}z'^3 Y^2 + \Phi'(z', w)Y + \Psi'(z', w) \quad (\text{Case II-A-II'-ii-i}).$$

Taking the coordinate change $Y' := Y + c'_{04}w^4$, we have

$$f_\lambda = x^2 + Y'^3 - \left(\frac{9}{2}c_{30}z'^3 + \frac{9}{2}c'_{21}z'^2 w + 3c'_{04}w^4\right)Y'^2 \\ + (9c_{30}c'_{04}z'^3 w^4 + \Phi'(z', w))Y' - \frac{9}{2}c_{30}c'^2_{04}z'^3 w^8 - c'_{04}w^4\Phi'(z', w) + \Psi'(z', w), \\ f_0 = x^2 - \left(\frac{9}{2}c'_{21}z'^2 w + 3c'_{04}w^4\right)Y'^2 - \frac{9}{2}c_{30}c'^2_{04}z'^3 w^8 - c'_{04}w^4\Phi'_0(z', w) + \Psi'_0(z', w).$$

Then $(f_0, 0)$ is an isolated singularity or Case II-A-I, and $1/2 + 17/50 + 3/25 + 2/25 > 1$, so $\{f_0 = 0\} - \{0\} \subset \mathbf{C}^4$ has at most rational singularities around the origin $0 \in \mathbf{C}^4$.

Remark 3.4. We will take the following arguments to show Theorem 2.7.

Taking suitable local coordinate changes finitely, if necessary, we have a three-dimensional face Δ of $\Gamma(f_\lambda)$ such that $(1, 1, 1, 1)$ lies strictly above the hyperplane including Δ , that $\text{Sing}(f_0) := \text{Sing}(\{f_0 = 0\}) = \bigcup_j C_j$ with $\dim_{\mathbf{C}} C_j \leq 1$, and that $(f_0, P) := (\{f_0 = 0\}, P)$ is rational for any $P \in C_j \subset \text{Sing}(f_0)$ with $P \neq 0$ and $0 \in C_j$, where $f_0 := f_\lambda \Delta = \sum_{\nu \in \Delta} a_\nu z^\nu \subset f_\lambda$. Therefore $\{f_0 = 0\} - \{0\}$ has at most rational singularities around 0, and so $(f_\lambda, 0)$ is rational.

More precisely, any three-dimensional face $\Delta = \{(X, Y, Z, W) \in \mathbf{R}_{\geq 0}^4; \frac{1}{2}X + \beta Y + \gamma Z + \delta W = 1, 0 \leq X \leq 2, j \leq Y \leq j', k \leq Z \leq k', l \leq W \leq l'\}$ of $\Gamma(f_\lambda)$ satisfies $1 < \frac{1}{2} + \beta + \gamma + \delta$ under the original local coordinates from Lemma 3.1. Choose a three-dimensional face Δ of $\Gamma(f_\lambda)$ suitably and let $f_0 := f_\lambda \Delta = \sum_{\nu \in \Delta} a_\nu z^\nu \subset f_\lambda := \sum_{\nu} a_\nu z^\nu$.

Case (0). $(f_0, 0)$ is an isolated singularity.

Then $(f_0, 0)$ is rational and so $\{f_0 = 0\} \subset \mathbf{C}^4$ has only rational singularities around the origin $0 \in \mathbf{C}^4$.

Case (I). $\text{Sing}(f_0) = \bigcup_j C_j$, $\dim_{\mathbf{C}} C_j \leq 1$ for all j .

Since f_0 is quasi-homogeneous, we have $\text{Sing}(C_j) = \{0\}$. If an arbitrary point $P \in C_j - \{0\}$ is rational on $\{f_0 = 0\}$ for each irreducible curve C_j with $0 \in C_j$, then $\{f_0 = 0\} - \{0\}$ has at most rational singularities around 0. Rationality of P is shown by using the following fact:

if there exists an hyperplane cut $(H, P) \subset (\{f_0 = 0\}, P)$ which is rational and Gorenstein, then $(\{f_0 = 0\}, P)$ is also rational and Gorenstein.

When we can not tell whether P is rational or not, take suitable local coordinate change and repeat the same procedure as above. The condition $1 < \frac{1}{2} + \beta^{(n)} + \gamma^{(n)} + \delta^{(n)}$ is still satisfied for a certain $\Delta^{(n)}$ after each coordinate change. This procedure must finish in finite times from the assumption $\mu(f_\lambda, 0) < \mu(f, 0)$.

Case (II). $\text{Sing}(f_0) = \bigcup_j C_j$, $\dim_{\mathbf{C}} C_j = 2$ for some j .

After suitable local coordinate change, choose another three-dimensional face Δ of $\Gamma(f_\lambda)$ properly, and let $f_0 := f_\lambda \Delta = \sum_{\nu \in \Delta} a_\nu z^\nu$.

(II-I⁽⁰⁾). $\text{Sing}(f_0) = \bigcup_j C_j$, $\dim_{\mathbf{C}} C_j \leq 1$ for all j .

Then the proof is completed similarly as in (I).

(II-II^(l)). $\text{Sing}(f_0) = \bigcup_j C_j$, $\dim_{\mathbf{C}} C_j = 2$ for some j .

After suitable local coordinates change, the condition $1 < \frac{1}{2} + \beta' + \gamma' + \delta'$ is still satisfied for a certain face $\Delta' = \{(X, Y, Z, W) ; \frac{1}{2}X + \beta'Y + \gamma'Z + \delta'W = 1\}$ of $\Gamma(f_\lambda)$. Choose such three-dimensional face Δ' properly; if Case (I^(l)) then the assertion is concluded. If Case (II^(l)) again, take suitable coordinate change once more. This procedure must finish in finite times from the assumption $\mu(f_\lambda, 0) < \mu(f, 0)$.

Proof of Theorem 2.7.

If there exists $y^j z^k w^l \in f_\lambda$ such that $j + k + l \leq 2$ then $(f_\lambda, 0)$ is at most rational. So we may assume $j + k + l \geq 3$ for all $y^j z^k w^l \in f_\lambda$.

Let $\Lambda := \{(k, l) ; z^k w^l \in f_\lambda\} \cup \{\frac{3}{2}(k, l) ; yz^k w^l \in f_\lambda\}$ and let Γ be the union of the compact faces of the convex hull of $\bigcup_{\nu \in \Lambda} (\nu + \mathbf{R}_{\geq 0}^2)$ in $\mathbf{R}^2$. For any one-dimensional face $\Delta = \{(Z, W) ; \gamma Z + \delta W = 1, k_1 \leq Z \leq k_0, l_0 \leq W \leq l_1\}$ of Γ , we get $\gamma + \delta > 1/6$ from Lemma 3.1. Choose such Δ satisfies $k_1 < 6$ and $l_0 < 6$. (See FIGURE 1.) Then,

$$f_\lambda = x^2 + y^3 + \sum_{\gamma k + \delta l \geq 2/3} b_{kl} y z^k w^l + \sum_{\gamma k + \delta l \geq 1} c_{kl} z^k w^l,$$

$$f_0 = x^2 + y^3 + \sum_{\gamma k + \delta l = 2/3} b_{kl} y z^k w^l + \sum_{\gamma k + \delta l = 1} c_{kl} z^k w^l$$

with respect to the weight $\alpha := (1/2, 1/3, \gamma, \delta)$.




FIGURE 1.

Case I. $h := f_0 - x^2 \in \mathbf{C}[y, z, w]$ has no double factor.

Since h and $\frac{\partial h}{\partial y}$ have no common factor, $\dim_{\mathbf{C}} \text{Sing}(f_0) \leq 1$. Let $C_j \subset \text{Sing}(f_0)$ be an irreducible curve with $0 \in C_j$. If $C_j \ni P = (0, a, b, c) \neq (0, 0, 0, 0)$ then $b \neq 0$ or $c \neq 0$. Let $P \neq 0$ be an arbitrary point on C_j .

(I-A). $C_j \ni P = (0, a(t), b(t), t) ; t \neq 0$.

Since f_0 is quasi-homogeneous, we have $a(t) = a't^{1/3\delta}$, $b(t) = b't^{\gamma/\delta}$ for some $a', b' \in \mathbf{C}$.

Let $\eta := y - a$, $\zeta := z - b$, and $f_0(w = t) := f_0(x, \eta + a, \zeta + b, t)$.

Then $f_0(w = t) - x^2 = \eta^3 + 3a\eta^2 + \dots \in \mathbf{C}[\eta, \zeta]$ has no double factor.

(In fact, if $f_0(w = t) - x^2 = (\eta + \varphi(\zeta))^2(\eta + \psi(\zeta))$ then

$$\begin{aligned} f_0(w = t) - x^2 &= \eta^3 + 3a\eta^2 - 3\varphi(\varphi - 2a)\eta - \varphi^2(2\varphi - 3a) \\ &= y^3 - 3(\varphi(z - b(t)) - a(t))^2y - 2(\varphi(z - b(t)) - a(t))^3, \\ f_0 - x^2 &= (y + \varphi(z - b(w)) - a(w))^2(y - 2\varphi(z - b(w)) + 2a(w)) \\ &= y^3 - 3(\varphi(z - b(w)) - a(w))^2y - 2(\varphi(z - b(w)) - a(w))^3. \end{aligned}$$

It follows that $\varphi(z - b(w)) - a(w) \in \mathbf{C}[z, w]$ from $\varphi(z - b(w)) - a(w) \in \mathbf{C}[z, w^{\gamma/\delta}]$, $(\varphi(z - b(w)) - a(w))^2, (\varphi(z - b(w)) - a(w))^3 \in \mathbf{C}[z, w]$.)

Therefore $(f_0(w = t), (0, 0, 0))$ is an isolated singularity under the local coordinate system (x, η, ζ) .

If $a \neq 0$ then $(f_0(x, y, z, t), (0, a, b))$ is rational, so $(f_0, (0, a, b, t))$ is rational.

If $a = 0$ then $\eta = y$, $f_0(w = t) = x^2 + y^3 + \sum b_{kl}y(\zeta + b)^kt^l + \sum c_{kl}(\zeta + b)^kt^l$.

If $b = 0$ then $\zeta = z$, $f_0(w = t) = x^2 + y^3 + \sum b_{kl}yz^kt^l + \sum c_{kl}z^kt^l$. By Lemma 3.2, $(f_0(x, y, z, t), (0, 0, 0))$ is rational, so $(f_0, (0, 0, 0, t))$ is rational.

So we consider the case $b \neq 0$.

If there exists $i \leq 5$ such that $\zeta^i \in f_0(w = t)$ or $j \leq 3$ such that $y\zeta^j \in f_0(w = t)$, then $(f_0(w = t), (0, 0, 0))$ is rational under the local coordinate system (x, y, ζ) . So we assume the coefficient of ζ^i is 0 for all $i \leq 5$ and the coefficient of $y\zeta^j$ is 0 for all $j \leq 3$.

Furthermore we may assume $\gamma \geq \delta$. (Indeed, if $\gamma < \delta$ we consider $f_0(x, y, s, w)$ instead of $f_0(x, y, z, t)$.) Thus $b = b(t)$ is written as $b(t) = b't^m$, $m := q/p = \gamma/\delta \geq 1$, $p, q \in \mathbf{N}$, $(p, q) = 1$.

Since the coefficient of ζ^i is $\frac{1}{i!} \frac{\partial^i f_0}{(\partial z)^i}(0, 0, b, t)$ and the coefficient of $y\zeta^j$ is $\frac{1}{j!} \frac{\partial^{j+1} f_0}{\partial y (\partial z)^j}(0, 0, b, t)$, it follows that:
the coefficient of ζ^i in $f_0(w = t)$ is 0

$$\begin{aligned} &\iff f_0(0, 0, z, t) = (z - b)^{i+1} \varphi(z, t), \text{ for some } \varphi(z, t) \in \mathbf{C}[z, t^m] \\ &\iff f_0(0, 0, z, w) = (z - b'w^m)^{i+1} \varphi(z, w), \text{ for some } \varphi(z, w) \in \mathbf{C}[z, w^m], \end{aligned}$$

and

$$\begin{aligned} &\text{the coefficient of } y\zeta^j \text{ in } f_0(w = t) \text{ is } 0 \\ &\iff \frac{\partial f_0}{\partial y}(0, 0, z, t) = (z - b)^{j+1} \psi(z, t), \text{ for some } \psi(z, t) \in \mathbf{C}[z, t^m] \\ &\iff \frac{\partial f_0}{\partial y}(0, 0, z, w) = (z - b'w^m)^{j+1} \psi(z, w), \text{ for some } \psi(z, w) \in \mathbf{C}[z, w^m]. \end{aligned}$$

The number $m = \gamma/\delta$ is a integer by the assumptions.

(In fact, if $p \neq 1$ then $f_0(0, 0, z, w)$ and $\frac{\partial f_0}{\partial y}(0, 0, z, w)$ are written as
 $f_0(0, 0, z, w) = (z - b'w^m)^6 \varphi(z, w) = (z^p - b''w^q)^6 \varphi'(z, w)$, $b'' \in \mathbf{C}$, $\varphi'(z, w) \in \mathbf{C}[z, w]$,
 $\frac{\partial f_0}{\partial y}(0, 0, z, w) = (z - b'w^m)^4 \psi(z, w) = (z^p - b'''w^q)^4 \psi'(z, w)$,

respectively. Since $2 \leq p \leq q$, we have $\frac{1}{6p} + \frac{1}{6q} \leq \frac{1}{6}$ and $\frac{1}{4p} + \frac{1}{4q} \leq \frac{1}{4}$. This is a contradiction to the condition $\gamma + \delta > 1/6$.)

Let $z' := z - b'w^m$, $\Lambda' := \{(k, l) ; z'^k w^l \in f_\lambda\} \cup \{\frac{3}{2}(k, l) ; yz'^k w^l \in f_\lambda\}$, and let Γ' be the union of the compact faces of the convex hull of $\bigcup_{\nu \in \Lambda'} (\nu + \mathbf{R}_{\geq 0}^2)$ in $\mathbf{R}^2$. Then:

Claim 3.5. For any one-dimensional face $\Delta' = \{\gamma'Z + \delta'W = 1, k'_1 \leq Z \leq k'_0, l'_0 \leq W \leq l'_1\}$ of Γ' , the condition $\gamma' + \delta' > 1/6$ is satisfied.

A proof of this claim is found at the end of this paper.

There exists $k_1 < i \leq k_0$ such that $i = \max\{i' \in \mathbf{N} ; (z - b'w^m)^{i'} | f_0(0, 0, z, w)\}$ or $\frac{2}{3}k_1 < j \leq \frac{2}{3}k_0$ such that $j = \max\left\{j' \in \mathbf{N} ; (z - b'w^m)^{j'} \left| \frac{\partial f_0}{\partial y}(0, 0, z, w) \right.\right\}$. Choose Δ' such that $k'_0 \leq \min\{i, \frac{3}{2}j\}$, $k'_1 < 6$ and $l'_0 < 6$. Then $k'_0 \leq k_0$, $l_0 \leq l'_0$, and $1 \leq \frac{\gamma}{\delta} < \frac{\gamma'}{\delta'}$. Let f_0 be the initial part of $f \in \mathbf{C}[x, y, z', w]$ with respect to the weight $(1/2, 1/3, \gamma', \delta')$. If $h := f_0 - x^2 \in \mathbf{C}[y, z', w]$ has a double factor, then Case (II-A). Now we assume h has no double factor. If $k'_0 < 6$ then our assertion is concluded. So we assume $k'_0 \geq 6$. Repeating same argument as above,

$$\begin{aligned} z'' &:= z' - b''w^{m'}, \dots, z^{(n)} := z^{(n-1)} - b^{(n)}w^{m^{(n-1)}}, \dots, \\ k_0 &\geq k'_0 \geq k''_0 \geq \dots \geq k_0^{(n)} \geq \dots, \\ 1 &\leq \frac{\gamma}{\delta} < \frac{\gamma'}{\delta'} < \frac{\gamma''}{\delta''} < \dots < \frac{\gamma^{(n)}}{\delta^{(n)}} < \dots, \\ \frac{1}{6} &< \gamma^{(n)} + \delta^{(n)} \text{ for all } n \in \mathbf{N}. \end{aligned}$$

(See FIGURE 2.) If there exists $n \in \mathbf{N}$ such that $k_0^{(n)} < 6$, then this proof is

completed. So consider the case $k_0^{(n)} \geq 6$ for all $n \in \mathbf{N}$. Then $\frac{\gamma^{(n)}}{\delta^{(n)}} \in \mathbf{N}$ for all $n \in \mathbf{N}$. If $k_0^{(n)} > 6$ for all $n \in \mathbf{N}$ then $\frac{\gamma^{(n)}}{\delta^{(n)}} < -\frac{l_0^{(n)} - 6}{k_0^{(n)} - 6}$, and so this procedure must finish in finite times (See FIGURE 3). Thus assume there exists $n \in \mathbf{N}$ such that $6 = k_0^{(n)} = k_0^{(n+1)} = k_0^{(n+2)} = \dots$. If this procedure continues infinitely then $\mu(f_\lambda, 0) \gg 1$, a contradiction. This completes Case (I-A).




FIGURE 2.

(I-B). $C_j \ni P = (0, a(s), s, c(s))$; $s \neq 0$.

Let $\eta := y - a$, $\omega := w - c$, and $f_0(z = s) := f_0(x, \eta + a, s, \omega + c)$. Then $f_0(z = s) - x^2 \in \mathbf{C}[\eta, \omega]$ has no double factor. If $a \neq 0$ or $a = c = 0$ then $(f_0(x, y, s, w), (0, a, c))$ is rational, so $(f_0, (0, a, s, c))$ is rational similarly as in Case (I-A). Thus we assume $a = 0$, $c \neq 0$. If $\gamma \geq \delta$ then our argument can be reduced to the case $a = 0$, $b \neq 0$ and $\gamma \geq \delta$ of Case (I-A). So it is sufficient to consider the case $\gamma < \delta$. Furthermore we may assume that the coefficient of ω^i in $f_0(z = s)$ is 0 for all $i \leq 5$ and the coefficient of $y\omega^j$ in $f_0(z = s)$ is 0 for all $j \leq 3$. Thus $c = c(s)$ is written as $c(s) = c's^m$, $c' \in \mathbf{C}$, $m = \delta/\gamma > 1$, $m \in \mathbf{N}$. Therefore $l_1 < 6$ for No.11-14, 46-51 and $l_1 \leq 6$ for No.10, 83. (Because $z^{12} \in f$ for No.10, $z^{10} \in f$ for No.11, $z^9 \in f$ for No.12, $z^8 \in f$ for No.13, $z^7 \in f$ for No.14, $z^{11} \in f$ for No.46, $yz^7 \in f$ for No.47, $z^9w \in f$ for No.48, $z^8w \in f$ for No.49, $z^7w^2 \in f$ for No.50, $z^7w \in f$ for No.51, $z^{10}w \in f$ for




FIGURE 3.

No.83.) Thus it is sufficient to consider the case $(k_1, l_1) = (0, 6)$ of No.10 and 83. Let $w' := w - c'z^m$, $\Lambda' := \{(k, l) ; z^k w^l \in f_\lambda\} \cup \{\frac{3}{2}(k, l) ; yz^k w^l \in f_\lambda\}$, and let Γ' be the union of the compact faces of the convex hull of $\bigcup_{\nu \in \Lambda'} (\nu + \mathbf{R}_{\geq 0}^2)$ in $\mathbf{R}^2$. Then for any one-dimensional face $\Delta' = \{\gamma'Z + \delta'W = 1, k'_1 \leq Z \leq k'_0, l'_0 \leq W \leq l'_1\}$ of Γ' , $\gamma' + \delta' > 1/6$ and $1 < \frac{\delta}{\gamma} < \frac{\delta'}{\gamma'}$ are satisfied. Repeating same argument as in Case (I-A),

$$w'' := w' - c''z^{m'}, \dots, w^{(n)} := w^{(n-1)} - c^{(n)}z^{m^{(n-1)}}, \dots,$$

$$l_1 \geq l'_1 \geq l''_1 \geq \dots \geq l_1^{(n)} \geq \dots,$$

$$1 < \frac{\delta}{\gamma} < \frac{\delta'}{\gamma'} < \frac{\delta''}{\gamma''} < \dots < \frac{\delta^{(n)}}{\gamma^{(n)}} < \dots,$$

$$\frac{1}{6} < \gamma^{(n)} + \delta^{(n)} \text{ for all } n \in \mathbf{N}.$$

If there exists $n \in \mathbf{N}$ such that $l_1^{(n)} < 6$ then the assertion holds true. If $l_1^{(n)} = 6$ for all $n \in \mathbf{N}$ then this procedure continues infinitely, therefore $\mu(f_\lambda, 0) \gg 1$, a contradiction. This completes Case (I).

For convenience, we write z, γ, δ , instead of $z^{(n)}, \gamma^{(n)}, \delta^{(n)}$, etc.

Case II. $h := f_0 - x^2 \in \mathbf{C}[y, z, w]$ has a double factor.

$$f_0 = x^2 + y^3 - \frac{27}{4}g(z, w)^2y - \frac{27}{4}g(z, w)^3 = x^2 + (y + \frac{3}{2}g(z, w))^2(y - 3g(z, w)).$$

Let $Y := y + \frac{3}{2}g(z, w)$, then f_0 and f_λ are written as:

$$f_0 = x^2 + Y^3 - \frac{9}{2}g(z, w)Y^2,$$

$$f_\lambda = x^2 + Y^3 - \frac{9}{2}g(z, w)Y^2 + \varphi(z, w)Y + \psi(z, w),$$

for some $\varphi(z, w), \psi(z, w) \in \mathbf{C}[z, w]$ with $\varphi \not\equiv 0$ or $\psi \not\equiv 0$, $\varphi \equiv 0$ or $\deg_\alpha \varphi > 2/3$, $\psi \equiv 0$ or $\deg_\alpha \psi > 1$.

(II-A). When $\gamma \geq \delta$, $g(z, w)$ can be classified into ten cases as below:

$$(1) \quad g = c_{10}z + c_{0L}w^L,$$

$$(2) \quad g = c_{11}zw + c_{0(L+1)}w^{L+1},$$

$$(3) \quad g = c_{20}z^2 + c_{0L}w^L, \quad 3 \leq L, \quad L \text{ is odd},$$

$$(4\text{-a}) \quad g = c_{20}(z + \gamma_1w^L)(z + \gamma_2w^L), \quad \gamma_1 \neq \gamma_2,$$

$$(4\text{-b}) \quad g = c_{20}(z + \gamma_1w^L)^2,$$

$$(5) \quad g = c_{21}z^2w + c_{0(L+1)}w^{L+1}, \quad 3 \leq L, \quad L \text{ is odd},$$

$$(6\text{-a}) \quad g = c_{21}(z + \gamma_1w^L)(z + \gamma_2w^L)w, \quad \gamma_1 \neq \gamma_2,$$

$$(6\text{-b}) \quad g = c_{21}(z + \gamma_1w^L)^2w,$$

$$(7) \quad g = c_{30}z^3 + c_{04}w^4,$$

$$(8) \quad g = c_{30}z^3 + c_{13}zw^3,$$

$$(9) \quad g = c_{30}z^3 + c_{05}w^5,$$

$$(10\text{-a}) \quad g = c_{30}(z + \gamma_1w)(z + \gamma_2w)(z + \gamma_3w), \quad \gamma_i \neq \gamma_j \text{ for } i \neq j,$$

$$(10\text{-b}) \quad g = c_{30}(z + \gamma_1w)^2(z + \gamma_2w), \quad \gamma_1 \neq \gamma_2,$$

$$(10\text{-c}) \quad g = c_{30}(z + \gamma_1w)^3.$$

After the local coordinate change $z' := z + \gamma_1w^L$ ($L = 1$ for (10)) around 0,

$$(4\text{-a}) \quad g = c_{20}z'^2 + c'_{1L}z'w^L,$$

$$(4\text{-b}) \quad g = c_{20}z'^2,$$

$$(6\text{-a}) \quad g = c_{21}z'^2w + c'_{1(L+1)}z'w^{L+1},$$

$$(6\text{-b}) \quad g = c_{21}z'^2w,$$

$$(10\text{-a}) \quad g = c_{30}z'^3 + c'_{21}z'^2w + c'_{12}z'w^2,$$

$$(10\text{-b}) \quad g = c_{30}z'^3 + c'_{21}z'^2w,$$

$$(10\text{-c}) \quad g = c_{30}z'^3.$$

(See FIGURE 4.)

Let $\Lambda' := \{(k, l) ; z'^kw^l \in f_\lambda\} \cup \{\frac{3}{2}(k, l) ; Yz'^kw^l \in f_\lambda\} \cup \{3(k, l) ; Y^2z'^kw^l \in f_\lambda\}$, and let Γ' be the union of the compact faces of the convex hull of $\bigcup_{\nu \in \Lambda'}(\nu + \mathbf{R}_{\geq 0}^2)$ in $\mathbf{R}^2$. Then for any one-dimensional face $\Delta' = \{\gamma'Z + \delta'W = 1, k'_1 \leq Z \leq k'_0, l'_0 \leq W \leq l'_1\}$




FIGURE 4.

of Γ' , the condition $\gamma' + \delta' > 1/6$ is satisfied. (In fact, No.11, 13, 14 and 49-51 can not become Case (10). About Case (10-c) which comes from No.10, 12, 46-48 or 83, the condition $\gamma' + \delta' > 1/6$ is satisfied as in Claim 3.5.)

First we consider Case (1), (2), (3), (4-a), (5), (6-a), (7), (8), (9) and (10-a).

We replace the weight $\alpha = (1/2, 1/3, \gamma, \delta)$ with $\alpha' = (1/2, \beta', \gamma', \delta')$ which satisfies the conditions

$$\begin{aligned} \frac{1}{3} = \beta < \beta' < \frac{1}{2} < \beta' + \gamma' + \delta', \quad \gamma'/\delta' = \gamma/\delta, \\ \deg_{\alpha'}(g(z, w)Y^2) = 1, \quad \deg_{\alpha'}(\varphi(z, w)Y), \deg_{\alpha'}\psi(z, w) \geq 1, \quad \text{and} \\ \deg_{\alpha'}(\varphi(z, w)Y) \text{ or } \deg_{\alpha'}\psi(z, w) = 1, \end{aligned}$$

and let f_0 be the initial part of f_λ . (There exists such weight α' because

$$\begin{aligned} \beta' + \gamma' + \delta' = \beta' + (\gamma + \delta)(1 - 2\beta')/(1 - 2\beta) > \frac{1}{2} \text{ and} \\ \deg_{\alpha'}(g(z, w)Y^2) = 2\beta' + \deg_\alpha g(z, w)(1 - 2\beta')/(1 - 2\beta) = 1 \end{aligned}$$

for all $\beta, \gamma, \delta, \beta', \gamma', \delta'$ satisfying

$$\beta, \beta' < \frac{1}{2} < \beta + \gamma + \delta \text{ and } \gamma'/\gamma = \delta'/\delta = (1 - 2\beta')/(1 - 2\beta).$$

Then

$$\begin{aligned} f_0 &= x^2 - \frac{9}{2}g(z, w)Y^2 + \varphi_0(z, w)Y + \psi_0(z, w), \\ f_\lambda &= x^2 + Y^3 - \frac{9}{2}g(z, w)Y^2 + \varphi(z, w)Y + \psi(z, w), \end{aligned}$$

where φ_0 (resp. ψ_0) is either 0 or the initial part of φ (resp. ψ).

(II-A-I). $h := f_0 - x^2 \in \mathbf{C}[Y, z, w]$ has no double factor.

Let $C_j \subset \text{Sing}(f_0)$ be an irreducible curve with $0 \in C_j$, and $P = (0, a, b, c) \neq 0$ an arbitrary point on C_j .

(II-A-I-i). $C_j \ni P = (0, a(t), b(t), t) ; t \neq 0$.

Then, $a(t) = a't^{\beta'/\delta'}$, $b(t) = b't^{\gamma'/\delta'}$ for some $a', b' \in \mathbf{C}$.

Let $\eta := Y - a$, $\zeta := z - b$, and $f_0(w = t) := f_0(x, \eta + a, \zeta + b, t)$, then

$$\begin{aligned} f_0(w = t) &= x^2 - \frac{9}{2}g(\zeta + b, t)(\eta + a)^2 + \varphi_0(\zeta + b, t)(\eta + a) + \psi_0(\zeta + b, t) \\ &= x^2 - \frac{9}{2}g(\zeta + b, t)\eta^2 + (-9ag(\zeta + b, t) + \varphi_0(\zeta + b, t))\eta \\ &\quad - \frac{9}{2}a^2g(\zeta + b, t) + a\varphi_0(\zeta + b, t) + \psi_0(\zeta + b, t). \end{aligned}$$

Then both $g(\zeta + b, t) \in \mathbf{C}[\zeta]$ and $f_0(w = t) - x^2 \in \mathbf{C}[\eta, \zeta]$ have no double factor.

Therefore $(f_0(w = t), (0, 0, 0))$ is an isolated singularity under the coordinates (x, η, ζ) .

If $g(b, t) \neq 0$ then $\eta^2 \in f_0(w = t)$. Otherwise, $\zeta \in g(\zeta + b, t)$, so $\eta^2\zeta \in f_0(w = t)$.

Hence $(f_0(w = t), (0, 0, 0))$ is rational under the coordinates (x, η, ζ) by Lemma 3.2.

(II-A-I-ii). $C_j \ni P = (0, a(s), s, 0) ; s \neq 0$.

Let $\eta := Y - a$, then

$$\begin{aligned} f_0(z = s) &= x^2 - \frac{9}{2}g(s, w)(\eta + a)^2 + \varphi_0(s, w)(\eta + a) + \psi_0(s, w) \\ &= x^2 - \frac{9}{2}g(s, w)\eta^2 + (-9ag(s, w) + \varphi_0(s, w))\eta \\ &\quad - \frac{9}{2}a^2g(s, w) + a\varphi_0(s, w) + \psi_0(s, w). \end{aligned}$$

$(f_0(z = s), (0, 0, 0))$ is an isolated singularity under the coordinates (x, η, w) because

$h := f_0(z = s) - x^2 \in \mathbf{C}[\eta, w]$ has no double factor. We have $\eta^2 \in f_0(z = s)$ for (1),

(3), (4-a), (7), (8), (9), (10-a), and $\eta^2w \in f_0(z = s)$ for (2), (5), (6-a), since $s \neq 0$.

Therefore $(f_0(z = s), (0, 0, 0))$ is rational.

(II-A-I-iii). $C_j \ni P = (0, a, 0, 0) ; a \neq 0$.

Since $f_0(Y = a) = x^2 - \frac{9}{2}g(z, w)a^2 + \varphi_0(z, w)a + \psi_0(z, w)$, it follows that

$(f_0(Y = a), (0, 0, 0))$ is rational under the coordinates (x, z, w) by Lemma 3.2.

(II-A-II). $h := f_0 - x^2 \in \mathbf{C}[Y, z, w]$ has a double factor.

$$\begin{aligned}
f_\lambda &= x^2 + Y^3 - \frac{9}{2}g(z, w)Y^2 + \varphi(z, w)Y + \psi(z, w), \\
f_0 &= x^2 - \frac{9}{2}g(z, w)Y^2 + \varphi_0(z, w)Y + \psi_0(z, w) \\
&= x^2 - \frac{9}{2}g(z, w)(Y + \phi(z, w))^2.
\end{aligned}$$

After the coordinate change $Y' := Y + \phi(z, w)$,

$$f_\lambda = x^2 + Y'^3 + (-\frac{9}{2}g - 3\phi)(z, w)Y'^2 + \varphi'(z, w)Y' + \psi'(z, w)$$

for some $\varphi', \psi' \in \mathbf{C}[z, w]$ with $\deg_{\alpha'} \varphi < \deg_{\alpha'} \varphi', \deg_{\alpha'} \psi < \deg_{\alpha'} \psi'$. We replace the weight $\alpha' = (1/2, \beta', \gamma', \delta')$ with $\alpha'' = (1/2, \beta'', \gamma'', \delta'')$ which satisfies the conditions

$$\begin{aligned}
\frac{1}{3} &< \beta'' < \frac{1}{2} < \beta'' + \gamma'' + \delta'', \quad \gamma''/\delta'' = \gamma'/\delta', \\
\deg_{\alpha''}(g(z, w)Y'^2) &= 1, \quad \deg_{\alpha''}(\varphi'(z, w)Y'), \deg_{\alpha''}\psi'(z, w) \geq 1, \quad \text{and} \\
\deg_{\alpha''}(\varphi'(z, w)Y') \text{ or } \deg_{\alpha''}\psi'(z, w) &= 1,
\end{aligned}$$

and let f_0 be the initial part of f_λ . Then

$$\begin{aligned}
f_0 &= x^2 - \frac{9}{2}g(z, w)Y'^2 + \varphi'_0(z, w)Y' + \psi'_0(z, w), \\
\beta' &< \beta'', \quad \gamma' > \gamma'', \quad \delta' > \delta'', \\
\deg_{\alpha'} g(z, w) &< \deg_{\alpha'} \phi(z, w), \\
\deg_{\alpha'} \varphi_0(z, w) &< \deg_{\alpha'} \varphi'_0(z, w) \text{ or } \varphi'_0(z, w) \equiv 0, \\
\deg_{\alpha'} \psi_0(z, w) &< \deg_{\alpha'} \psi'_0(z, w) \text{ or } \psi'_0(z, w) \equiv 0.
\end{aligned}$$

If $h := f_0 - x^2 \in \mathbf{C}[Y', z, w]$ has no double factor, then Case (II-A-I).

If h has a double factor, namely,

$$f_0 = x^2 - \frac{9}{2}g(z, w)(Y' + \phi'(z, w))^2,$$

then after the coordinate change $Y'' := Y' + \phi'(z, w)$,

$$\begin{aligned}
f_\lambda &= x^2 + Y''^3 + (-\frac{9}{2}g - 3(\phi + \phi'))(z, w)Y''^2 + \varphi''(z, w)Y'' + \psi''(z, w) \\
&=: x^2 + Y''^3 + (-\frac{9}{2}g - 3\phi)(z, w)Y''^2 + \varphi''(z, w)Y'' + \psi''(z, w)
\end{aligned}$$

for some $\varphi'', \psi'' \in \mathbf{C}[z, w]$ with $\deg_{\alpha''} \varphi' < \deg_{\alpha''} \varphi'', \deg_{\alpha''} \psi' < \deg_{\alpha''} \psi''$.

If this procedure continues infinitely, then

$$\begin{aligned}
f_\lambda &= x^2 + Y^{(n)3} + (-\frac{9}{2}g - 3\phi)(z, w)Y^{(n)2} + \varphi^{(n)}(z, w)Y^{(n)} + \psi^{(n)}(z, w), \\
\frac{1}{3} &= \beta < \beta' < \beta'' < \dots < \beta^{(n)} < \dots < \frac{1}{2}, \\
\gamma &> \gamma' > \gamma'' > \dots > \gamma^{(n)} > \dots, \\
\delta &> \delta' > \delta'' > \dots > \delta^{(n)} > \dots, \\
\gamma/\delta &= \gamma'/\delta' = \gamma''/\delta'' = \dots = \gamma^{(n)}/\delta^{(n)} = \dots, \\
\frac{1}{2} &< \beta^{(n)} + \gamma^{(n)} + \delta^{(n)} \quad \text{for all } n \in \mathbf{Z}_{\geq 0}, \\
\deg_\alpha g(z, w) &< \deg_\alpha \phi(z, w), \\
\deg_\alpha \varphi_0(z, w) &< \deg_\alpha \varphi'_0(z, w) < \deg_\alpha \varphi''_0(z, w) < \dots < \deg_\alpha \varphi_0^{(n)}(z, w) < \dots,
\end{aligned}$$

$\deg_\alpha \psi_0(z, w) < \deg_\alpha \psi'_0(z, w) < \deg_\alpha \psi''_0(z, w) < \cdots < \deg_\alpha \psi_0^{(n)}(z, w) < \cdots$,
so $\mu(f_\lambda, 0) \geq \dim_{\mathbf{C}} \mathbf{C}\{w\} \Big/ \left(\varphi^{(n)}(0, w), \frac{\partial \psi^{(n)}}{\partial z}(0, w), \frac{\partial \psi^{(n)}}{\partial w}(0, w) \right) \gg 1$,
a contradiction.

Next we consider Case (4-b), (6-b), (10-b), (10-c). Choose a Δ' such that:

$$(k'_0, l'_0) = (6, 0) \text{ for (4-b), } (6, 3) \text{ for (6-b) and (10-b),}$$

$$k'_1 < 6 \text{ and } l'_0 < 6 \text{ for (10-c),}$$

respectively. Then $1 \leq \frac{\gamma}{\delta} < \frac{\gamma'}{\delta'}$. Let f_0 be the initial part of $f_\lambda \in \mathbf{C}[x, Y, z', w]$ with respect to the weight $\alpha' := (1/2, 1/3, \gamma', \delta')$, and $m' := \gamma'/\delta'$.

For the case $z'^3 Y^2$ is not contained in f_0 of (10-c), the situation is similar to (I-A), but $h := f_0 - x^2 \in \mathbf{C}[Y, z^{(n)}, w]$ can not have a double factor for $n \in \mathbf{N}$. So we consider other cases, i.e.,

$$f_0 = x^2 + Y^3 - \frac{9}{2} c_{kl}^{(n)} z'^k w^l Y^2 + \varphi'_0(z', w)Y + \psi'_0(z', w)$$

$$= \begin{cases} x^2 + Y^3 - \frac{9}{2} c_{20} z'^2 Y^2 + \varphi'_0(z', w)Y + \psi'_0(z', w), & (4\text{-b}) \\ x^2 + Y^3 - \frac{9}{2} c_{21} z'^2 w Y^2 + \varphi'_0(z', w)Y + \psi'_0(z', w), & (6\text{-b}) \\ x^2 + Y^3 - \frac{9}{2} c'_{21} z'^2 w Y^2 + \varphi'_0(z', w)Y + \psi'_0(z', w), & (10\text{-b}) \\ x^2 + Y^3 - \frac{9}{2} c_{30} z'^3 Y^2 + \varphi'_0(z', w)Y + \psi'_0(z', w), & (10\text{-c}) \end{cases}$$

where $\deg \varphi'_0(z', t) < 2k$, $\deg \psi'_0(z', t) < 3k$ ($0 \neq t \in \mathbf{C}$).

(II-A-I'). $h := f_0 - x^2 \in \mathbf{C}[Y, z', w]$ has no double factor.

(II-A-I'-i). $\text{Sing}(f_0) \supset C_j \ni P = (0, a(t), b(t), t)$; $t \neq 0$.

Let $\eta := Y - a$, $\zeta := z' - b$ and $f_0(w = t) := f_0(x, \eta + a, \zeta + b, t)$, then $f_0(w = t) - x^2$ has no double factor.

(Indeed, assume $f_0(w = t) - x^2 = (\eta + A(\zeta))^2(\eta + B(\zeta))$ for some $A, B \in \mathbf{C}[\zeta]$, then

$$\begin{aligned} f_0(w = t) - x^2 &= (Y - a(t) + A(z' - b(t)))^2(Y - a(t) + B(z' - b(t))), \\ f_0 - x^2 &= (Y - a(w) + A(z' - b(w)))^2(Y - a(w) + B(z' - b(w))) \\ &= (Y + G(z', w))^2(Y + H(z', w)) \\ (G &:= -a(w) + A(z' - b(w)), H := -a(w) + B(z' - b(w)) \in \mathbf{C}[z', w^{m'}].) \\ &= Y^3 + (2G + H)Y^2 + G(G + 2H)Y + G^2H \\ &= Y^3 - \frac{9}{2}g'Y^2 - G(9g' + 3G)Y - G^2(\frac{9}{2}g' + 2G), \end{aligned}$$

where $g'(z', w) := c_{kl}^{(l)} z'^k w^l$. It follows that $\deg G(z', t) < k$ from $\deg \varphi'_0(z', t) < 2k$, $\deg \psi'_0(z', t) < 3k$, and that $G \in \mathbf{C}[z', w]$ from g' , $G(9g' + 3G)$, $G^2(\frac{9}{2}g' + 2G) \in \mathbf{C}[z', w]$, $G \in \mathbf{C}[z', w^{m'}]$.)

If $a = b = 0$ then

$$f_0(w = t) = x^2 + Y^3 - \frac{9}{2}c_{kl}^{(l)} z'^k t^l Y^2 + \varphi'_0(z', t)Y + \psi'_0(z', t),$$

with $\text{ord}\varphi'_0(z', t) < 4 \leq 2k$ or $\text{ord}\psi'_0(z', t) < 6 \leq 3k$.

If $a = 0$, $b \neq 0$, then

$$\begin{aligned} f_0(w = t) &= x^2 + Y^3 - \frac{9}{2}c_{kl}^{(l)} (\zeta^k + kb\zeta^{k-1} + \dots + kb^{k-1}\zeta + b^k) t^l Y^2 \\ &\quad + \varphi'_0(\zeta + b, t)Y + \psi'_0(\zeta + b, t) \ni Y^2. \end{aligned}$$

If $a \neq 0$, $b = 0$, then

$$f_0(w = t) = x^2 + \eta^3 + 3a\eta^2 + (3a^2 + \varphi'_0(z', t))\eta - \frac{9}{2}c_{kl}^{(l)} z'^k t^l (\eta + a)^2 + \dots \ni \eta^2.$$

If $a \neq 0$, $b \neq 0$, then $\eta^2 \in f_0(w = t)$.

(Indeed, the coefficient of η^2 in $f_0(w = t)$ is $\frac{1}{2} \frac{\partial^2 f_0}{(\partial Y)^2}(0, a, b, t)$, so it follows that:

the coefficient of η^2 in $f_0(w = t)$ is 0

$$\iff f_0(0, Y, b(t), t) = (Y - a(t))^3$$

$$\iff f_0(0, Y, b(w), w) = (Y - a(w))^3$$

$$\iff f_0(0, Y, z', w) = (Y - \frac{3}{2}c_{kl}^{(l)} z'^k w^l)^3.)$$

Hence $(f_0(w = t), 0)$ is rational.

(II-A-I'-ii). $\text{Sing}(f_0) \supset C_j \ni P = (0, a(s), s, 0)$; $s \neq 0$.

Then $a = 0$ because $0 = f_0(0, a, s, 0) = \frac{\partial f_0}{\partial Y}(0, a, s, 0) = \varphi'_0(s, 0) = \psi'_0(s, 0)$.

Furthermore,

$$f_0(z' = s) - x^2 := f_0(x, Y, s, w) - x^2 = Y^3 - \frac{9}{2}c_{kl}^{(l)} s^k w^l Y^2 + \varphi'_0(s, w)Y + \psi'_0(s, w)$$

has no double factor. So it follows that $(f_0(z' = s), 0)$ is rational from

$$Y^2 \in f_0(z' = s) \text{ for (4-b) and (10-c),}$$

$$Y^2 w \in f_0(z' = s) \text{ and } w^i, Yw^j \text{ are not contained in } f_0(z' = s) \text{ for } i \leq 3, j \leq 2$$

$$\text{for (6-b) and (10-b).}$$

(II-A-II'). $h := f_0 - x^2 \in \mathbf{C}[Y, z', w]$ has a double factor.

f_0 and f_λ are written as:

$$\begin{aligned} f_0 &= x^2 + (Y + G(z', w))^2(Y + (-\frac{9}{2}g' - 2G)(z', w)) \\ &=: x^2 + Y'^3 - (\frac{9}{2}g' + 3G)(z', w)Y'^2, \quad (Y' := Y + G(z', w).) \end{aligned}$$

$$f_\lambda =: \begin{cases} x^2 + Y'^3 - \frac{9}{2}g''(z', w)Y'^2 + \varphi''(z', w)Y' + \psi''(z', w), & (4\text{-b}, 6\text{-b}, 10\text{-c}) \\ x^2 + Y'^3 - \frac{9}{2}(c_{30}z'^3 + g''(z', w))Y'^2 + \varphi''(z', w)Y' + \psi''(z', w), & (10\text{-b}) \end{cases}$$

for some $g'' := g' + \frac{2}{3}G$, $\varphi'', \psi'' \in \mathbf{C}[z', w]$ with $\deg G(z', t) < k$ ($0 \neq t \in \mathbf{C}$),

$\deg_{\alpha'} \varphi'' > 2/3 = 2(k\gamma' + l\delta')$, $\deg_{\alpha'} \psi'' > 1$.

Then g'' is one of the following:

$$\left. \begin{aligned} (3) \quad & g'' = c_{20}z'^2 + c'_{0L'}w^{L'}, \quad 3 \leq L', \quad L' \text{ is odd}, \\ (4\text{-a}) \quad & g'' = c_{20}(z' + \gamma'_1 w^{L'})(z' + \gamma'_2 w^{L'}), \quad 2 \leq L', \quad \gamma'_1 \neq \gamma'_2, \\ (4\text{-b}) \quad & g'' = c_{20}(z' + \gamma'_1 w^{L'})^2, \quad 2 \leq L', \\ (5) \quad & g'' = c_{21}^{(l)}z'^2 w + c'_{0(L'+1)}w^{L'+1}, \quad 3 \leq L', \quad L' \text{ is odd}, \\ (6\text{-a}) \quad & g'' = c_{21}^{(l)}(z' + \gamma'_1 w^{L'})(z' + \gamma'_2 w^{L'})w, \quad 2 \leq L', \quad \gamma'_1 \neq \gamma'_2, \\ (6\text{-b}) \quad & g'' = c_{21}^{(l)}(z' + \gamma'_1 w^{L'})^2 w, \quad 2 \leq L', \\ (7) \quad & g'' = c_{30}z'^3 + c'_{04}w^4, \\ (8) \quad & g'' = c_{30}z'^3 + c'_{13}z'w^3, \\ (9) \quad & g'' = c_{30}z'^3 + c'_{05}w^5, \end{aligned} \right\} \begin{aligned} & \text{(become from 4-b)} \\ & \text{(become from} \\ & \text{6-b or 10-b)} \\ & \text{(become from 10-c).} \end{aligned}$$

For 3, 4-a, 5, 6-a, 7, 8 and 9, we replace the weight $\alpha' = (1/2, 1/3, \gamma', \delta')$ with $\alpha'' = (1/2, \beta'', \gamma'', \delta'')$ which satisfies the conditions

$$\frac{1}{3} < \beta'' < \frac{1}{2} < \beta'' + \gamma'' + \delta'', \quad \gamma''/\delta'' = \gamma'/\delta',$$

$$\deg_{\alpha''}(g''(z', w)Y'^2) = 1, \quad \deg_{\alpha''}(\varphi''(z', w)Y'), \deg_{\alpha''}\psi''(z', w) \geq 1, \quad \text{and}$$

$$\deg_{\alpha''}(\varphi''(z', w)Y') \text{ or } \deg_{\alpha''}\psi''(z', w) = 1,$$

and let f_0 be the initial part of f_λ .

If $h := f_0 - x^2 \in \mathbf{C}[Y', z', w]$ has no double factor then Case (II-A-I). Otherwise, Case (II-A-II), and this procedure must finish in finite times from the assumption.

For 4-b, 6-b, after the coordinate change $z'' := z' + \gamma'_1 w^{L'}$, we define Λ'', Γ'' under the coordinates (x, Y', z'', w) similarly as before and choose a Δ'' corresponding to a weight $\alpha'' = (1/2, 1/3, \gamma'', \delta'')$ which satisfies $(k_0'', l_0'') = (6, 0)$ for (4-b), $(6, 3)$ for (6-b). Then $\frac{1}{6} < \gamma'' + \delta''$ and $1 \leq \frac{\gamma}{\delta} < \frac{\gamma'}{\delta'} < \frac{\gamma''}{\delta''}$. Let f_0 be the initial part of f_λ . Then,

$$f_0 = \begin{cases} x^2 + Y'^3 - \frac{9}{2}c_{20}z''^2Y'^2 + \varphi_0'''(z'', w)Y' + \psi_0'''(z'', w), & (4-b) \\ x^2 + Y'^3 - \frac{9}{2}c_{21}^{(t)}z''^2wY'^2 + \varphi_0'''(z'', w)Y' + \psi_0'''(z'', w), & (6-b) \end{cases}$$

where $\deg \varphi_0'''(z'', t) < 4$, $\deg \psi_0'''(z'', t) < 6$ ($0 \neq t \in \mathbf{C}$).

If $h := f_0 - x^2 \in \mathbf{C}[Y', z'', w]$ has no double factor then Case (II-A-I'). Otherwise, Case (II-A-II'), and this procedure must finish in finite times from the assumption.

(II-B). When $\gamma < \delta$, $g(z, w)$ can be classified as below:

- (1') $g = c_{01}w + c_{K0}z^K$, $2 \leq K$,
- (2') $g = c_{11}zw + c_{(K+1)0}z^{K+1}$, $2 \leq K$,
- (3') $g = c_{02}w^2 + c_{K0}z^K$, $3 \leq K$, K is odd,
- (4-a') $g = c_{02}(w + \gamma_1 z^K)(w + \gamma_2 z^K)$, $2 \leq K$, $\gamma_1 \neq \gamma_2$, (for No.10, 83 only),
- (4-b') $g = c_{02}(w + \gamma_1 z^K)^2$, $2 \leq K$, (for No.10, 83 only),
- (5') $g = c_{12}zw^2 + c_{(K+1)0}z^{K+1}$, $3 \leq K$, K is odd, (for No.10 only),
- (7') $g = c_{03}w^3 + c_{40}z^4$, (for No.10 only).

For Cases (1'), (2'), (3'), (4-a'), (5') and (7'), the situation is similar to (II-A). For Case (4-b') of No.10 and 83, we define $\Lambda', \Gamma', \Delta', f_0$ and h similarly as in (II-A), and repeat the same argument as in (II-A). Namely, if $h := f_0 - x^2 \in \mathbf{C}[Y, z, w']$ has no double factor then (II-B-I'), and if h has a double factor then (II-B-II'). Then the condition $\beta^{(n)} + \gamma^{(n)} + \delta^{(n)} > \frac{1}{2}$ must be always satisfied for No.10, because $\mu(f_\lambda, 0) < \mu(f, 0) = 242$. For the case of $(k_1, l_1) = (k'_1, l'_1) = (0, 6)$ of No.83, if it becomes (II-B-II'), then $z^k \in f_\lambda$ for some $k \geq 15$. And so, if $\beta^{(n)} + \gamma^{(n)} + \delta^{(n)} \leq 1/2$ for some $n \in \mathbf{N}$ then $\mu(f_\lambda, 0) \geq (3-1)(10-1)(15-1) = 252$, which is a contradiction to the condition $\mu(f_\lambda, 0) < \mu(f, 0) = 245$.

And this argument must finish finitely since $\mu(f_\lambda, 0) < \mu(f, 0)$.

For arrangement, let us illustrate the above argument as follows:

$$\text{I} \left\{ \begin{array}{l} \text{I-A} \left\{ \begin{array}{l} a \neq 0 \quad : \text{O.K.} \dots (*) \\ a = b = 0 : \text{O.K.} \dots (**) \\ a = 0, b \neq 0 \left\{ \begin{array}{l} \zeta^i (\exists i \leq 5) \text{ or } y\zeta^j (\exists j \leq 3) \in f_0(w = t) : \text{O.K.} \dots (***) \\ \text{otherwise} : z' := z - b'w^m \longrightarrow \left\{ \begin{array}{l} \text{I-A} \\ \text{II-A} \end{array} \right. \end{array} \right. \end{array} \right. \\ \\ \text{I-B} \left\{ \begin{array}{l} a \neq 0 \quad : \text{O.K.} \dots (*') \\ a = c = 0 : \text{O.K.} \dots (**') \\ a = 0, c \neq 0 \left\{ \begin{array}{l} \omega^i (\exists i \leq 5) \text{ or } y\omega^j (\exists j \leq 3) \in f_0(z = s) : \text{O.K.} \dots (***) \\ \text{otherwise} : w' := w - c'z^m \longrightarrow \left\{ \begin{array}{l} \text{I-B} \\ \text{II-B} \end{array} \right. \end{array} \right. \end{array} \right. \end{array} \right.$$

$$\begin{array}{l}
\text{II} \left\{ \begin{array}{l}
\begin{array}{l}
\text{II-A} \left\{ \begin{array}{l}
1, 2, 3, 4\text{-a}, 5, \\
6\text{-a}, 7, 8, 9, 10\text{-a}
\end{array} \rightarrow \begin{cases} \text{II-A-I : O.K.} \dots (*'') \\ \text{II-A-II : } Y' := Y + \phi \rightarrow \begin{cases} \text{II-A-I : O.K.} \dots (*'') \\ \text{II-A-II : } Y'' := Y' + \phi' \rightarrow \dots \end{cases} \end{cases} \\
\\
\begin{array}{l}
4\text{-b} \\
6\text{-b} : z' := z + \gamma_1 w^L \rightarrow \\
10\text{-b}
\end{array} \rightarrow \begin{cases} \text{II-A-I}' : \text{O.K.} \dots (**'') \\ \text{II-A-II}' : Y' := Y + G \rightarrow \begin{cases} 3, 4\text{-a}, \begin{cases} \text{II-A-I : O.K.} \dots (*'') \\ \text{II-A-II : } Y'' := Y' + \phi' \rightarrow \end{cases} \\ 4\text{-b} : z'' := z' + \gamma'_1 w^{L'} \\ 6\text{-b} \rightarrow \begin{cases} \text{II-A-I}' : \text{O.K.} \dots (**'') \\ \text{II-A-II}' : Y'' := Y' + G' \rightarrow \end{cases} \end{cases} \end{cases} \\
\\
10\text{-c} : z' := z + \gamma_1 w \rightarrow \begin{cases} z'^3 Y^2 \in f_0 \begin{cases} \text{II-A-I}' : \text{O.K.} \dots (**'') \\ \text{II-A-II}' : Y' := Y + G \rightarrow \begin{cases} \text{II-A-I : O.K.} \dots (*'') \\ \text{II-A-II : } Y'' := Y' + \phi' \rightarrow \end{cases} \\ \text{otherwise : I-A} \end{cases} \end{cases}
\end{array} \right. \\
\\
\text{II-B} \left\{ \begin{array}{l}
\begin{array}{l}
1', 2', 3', \\
4\text{-a}', 5', 7'
\end{array} \rightarrow \begin{cases} \text{II-B-I : O.K.} \dots (*''') \\ \text{II-B-II : } Y' := Y + \phi \rightarrow \begin{cases} \text{II-B-I : O.K.} \dots (*''') \\ \text{II-B-II : } Y'' := Y' + \phi' \rightarrow \dots \end{cases} \end{cases} \\
\\
4\text{-b}' : w' := w + \gamma_1 z^K \rightarrow \begin{cases} \text{II-B-I}' : \text{O.K.} \dots (**''') \\ \text{II-B-II}' : Y' := Y + G \rightarrow \begin{cases} 3', 4\text{-a}' \begin{cases} \text{II-B-I : O.K.} \dots (*''') \\ \text{II-B-II : } Y'' := Y' + \phi' \rightarrow \end{cases} \\ 4\text{-b}' : w'' := w' + \gamma'_1 z^{K'} \rightarrow \begin{cases} \text{II-B-I}' : \text{O.K.} \dots (**''') \\ \text{II-B-II}' : Y'' := Y' + G' \rightarrow \end{cases} \end{cases} \end{cases}
\end{array} \right.
\end{array}
\end{array}$$

The procedures of combinations of “a coordinate change $\longrightarrow$ ” must finish in finite times. Namely, taking suitable coordinate changes in finite times if necessary, the situation can be reduced to the case that $(\{f_0 = 0\} \cap H, P)$ is rational i.e. the case $(*\cdots*(^{(m)}))$ of (I) or (II), after all. Thus, there exist a local coordinate system and a weight such that (f_0, P) is rational. Furthermore, the condition $\frac{1}{2} + \beta^{(n)} + \gamma^{(n)} + \delta^{(n)} > 1$ is satisfied under each coordinate system appearing in the above procedures.

Thus it is enough to show Claim 3.5 for the proof of Theorem 2.7.

Proof of Claim 3.5.

Step 1.

Since $(f_\lambda, 0)$ is an isolated singularity, it follows that:

$$z^k \text{ or } yz^k \text{ or } z^kw \in f_\lambda, \quad \text{and} \quad w^l \text{ or } yw^l \text{ or } zw^l \in f_\lambda.$$

Taking suitable coordinate change

$$y' := y + b_1 z^{K_1} + c_1 w^{L_1}, \quad z' := z + c_2 w^{L_2}, \quad w' := w + b_2 z^{K_2}$$

for some sufficiently large $K_i, L_j \in \mathbf{N}$ and some $b_i, c_j \in \mathbf{C}$, we have $z'^{k'}, w'^{l'} \in f_\lambda$ and $\Gamma(f_\lambda)$ is the same as before except adding compact faces touching some coordinate planes.

Step 2 ([A]-Thm. XXII, [AGV]-12.7, Kouchnirenko [K]-Thm. I).

Let $y^J, z^K, w^L \in F(y, z, w) \in \mathbf{C}[y, z, w]$ and $V = \bigcup_{i=0}^3 V_i$ be a decomposition of the three-dimensional region of the positive orthant below the Newton boundary $\Gamma(F) \subset \mathbf{R}_{\geq 0}^3 = \{(Y, Z, W) \in \mathbf{R}_{\geq 0}^3\}$ which satisfies:

- V_0 is the three-dimensional simplicial cone which has vertex $(0, 0, 0)$, $(j, 0, 0)$, $(0, k, 0)$, $(0, 0, l)$ with $j, k, l \in \mathbf{Q}_{>0}$; $j \leq J, k \leq K, l \leq L$,
- V_1 has a vertex $(J, 0, 0)$, V_2 has a vertex $(0, K, 0)$, V_3 has a vertex $(0, 0, L)$,
- $\dim_{\mathbf{R}}(V_i \cap V_j) \leq 2$ for $i \neq j$, and $S_{12} = S_{23} = S_{31} = \emptyset$ for
- $S_{i1} := V_i \cap (YZ\text{-plane}), S_{i2} := V_i \cap (ZW\text{-plane}), S_{i3} := V_i \cap (WY\text{-plane}).$

(See FIGURE 5.)




FIGURE 5.

Suppose that $(F, 0)$ is an isolated singularity, then

$$\begin{aligned}
 \mu(F, 0) &\geq 3! \left(\sum_{i=0}^3 V_i \right) - 2! \left(\sum_{i=0}^3 \sum_{j=1}^3 S_{ij} \right) + 1!(J + K + L) - 1 \\
 &= jkl - (jk + kl + lj) + j + k + l - 1 \\
 &\quad + \sum_{i=1}^3 \left(6V_i - 2 \sum_{j=1}^3 S_{ij} \right) + (J - j) + (K - k) + (L - l) \\
 &\geq (j - 1)(k - 1)(l - 1).
 \end{aligned}$$

Step 3.

If $0 < \delta' < \delta \leq \gamma < \gamma' < 1$ and $\gamma + \delta = \gamma' + \delta'$ then

$$\left(\frac{1}{\gamma} - 1\right) \left(\frac{1}{\delta} - 1\right) < \left(\frac{1}{\gamma'} - 1\right) \left(\frac{1}{\delta'} - 1\right).$$

In fact, let $\gamma + \delta = \gamma' + \delta' =: 1/c$, $c \in \mathbf{R}$, then

$$\begin{aligned} & \frac{1}{\gamma'\delta'} - \frac{1}{\gamma'} - \frac{1}{\delta'} - \left(\frac{1}{\gamma\delta} - \frac{1}{\gamma} - \frac{1}{\delta}\right) \\ &= (c-1)(\gamma' - \gamma)(c(\gamma + \gamma') - 1)/\gamma\gamma'(1 - c\gamma)(1 - c\gamma') > 0. \end{aligned}$$

Step 4.

From Step 1 - Step 3, if $\gamma' + \delta' = 1/c \leq 1/6$ then

$$\begin{aligned} \mu(f_\lambda, 0) &\geq (3-1) \left(\frac{1}{\gamma'} - 1\right) \left(\frac{1}{\delta'} - 1\right) \\ &\geq (3-1) \left(\frac{6}{c\gamma'} - 1\right) \left(\frac{6}{c\delta'} - 1\right) \\ &> (3-1) \left(\frac{1}{\alpha_3} - 1\right) \left(\frac{1}{\alpha_4} - 1\right) = \mu(f, 0), \end{aligned}$$

a contradiction. This completes the proof of Claim 3.5 and Theorem 2.7. Q.E.D.

REFERENCES

- [A] V. I. ARNOL'D, *Critical points of smooth functions and their normal forms*, Uspekhi Mat. Nauk 30:5 (1975), 3-65. MR 54 #8701. = Russian Math. Surveys 30:5 (1975), 1-75.
- [AGV] V. I. ARNOLD, S. M. GUSEIN-ZADE, A. N. VARCHENKO, *Singularities of differentiable maps, I*, Monographs in Mathematics, Vol.83, Birkhäuser, Boston, Basel, Berlin, 1985.
- [E] R. ELKIK, *Singularites rationnelles et deformations*, Inventiones Math. 47 (1978), 139-147.
- [I1] S. ISHII, *On isolated Gorenstein singularities*, Math. Ann. 270 (1985), 541-554.
- [I2] ———, *The canonical modifications by weighted blow-ups*, to appear in Journal of Algebraic Geometry.
- [IW] S. ISHII, K. WATANABE, *A geometric characterization of a simple K3 singularity*, Tohoku Math.J. 44 (1992), 19-24.
- [K] A. G. KOUCHNIRENKO, *Polyèdres de Newton et nombres de Milnor*, Inventiones Math. 32 (1976), 1-31.
- [KKMS] G. KEMPF, F. KNUDSEN, D. MUMFORD, B. SAINT-DONAT, *Toroidal embeddings I*, Lecture Notes in Math., Springer, 339 (1973).

- [KS] A. KAS, M. SCHLESSINGER, *On the versal deformation of a complex space with an isolated singularity*, Math. Ann. 196 (1972), 23-29.
- [M] J. MILNOR, *Singular points of complex hypersurfaces*, Annals of Math. Studies 61, Princeton University Press, 1968.
- [R1] M. REID, *Canonical 3-folds*, Journées de Géométrie algébrique d'Angers, (A. Beauville, ed.), Sijthoff and Noordhoff, Alphen aan den Rijn, 1980, 273-310.
- [R2] ———, *Young person's guide to canonical singularities*, Proceedings of Symposia in Pure Mathematics, 46 (1987), 345-414.
- [Te] B. TEISSIER, *The hunting of invariants in the geometry of discriminants*, Real and Complex Singularities, Oslo, 1976, (P. Holm, ed.), Sijthoff and Noordhoff, (1977) 565-677.
- [Tj] G. N. TJURINA, *Locally semiuniversal flat deformations of isolated singularities of complex spaces*, Izv. Akad. Nauk SSSR, Ser. Mat. 33, No.5 (1969), 1026-1058 (In Russian).
- [TW] M. TOMARI, K-i. WATANABE, *Filtered rings, filtered blowing-ups and normal two-dimensional singularities with "star-shaped" resolution*, Publ. Res. Inst. Math. Sci., Kyoto Univ. 25 (1989), 681-740.
- [V1] A. N. VARCHENKO, *Zeta-Function of monodromy and Newton's diagram*, Invent. Math. 37 (1976), 253-262.
- [V2] ———, *A lower bound for the codimension of the stratum $\mu = \text{const}$ in term of the mixed hodge structure*, Vestnik Moskovskogo Universiteta. Matematika, Vol.37, No.6 (1982), 28-31.
- [W] K. WATANABE, *On plurigenera of normal isolated singularities II*, in Complex Analytic Singularities (T. Suwa, P. Wagreich, eds.), Advanced Studies in Pure Math. 8, Kinokuniya, Tokyo and North-Holland, Amsterdam, New York, Oxford, 1986, 671-685.
- [Ya] S. S. T. YAU, *Sheaf cohomology on 1-convex manifolds*, "Recent Developments in Several Complex Variables", Ann. of Math. Studies, Princeton Univ., 100 (1981), 429-452.
- [Yo] T. YONEMURA, *Hypersurface simple K3 singularities*, Tohoku Math.J. 42 (1990), 351-380.

E-mail address: furuya@math.titech.ac.jp